arXiv:math/9907175v2 [math.QA] 9 Dec 1999

QUANTUM VERTEX REPRESENTATIONS VIA FINITE GROUPS AND THE MCKAY CORRESPONDENCE

IGOR B. FRENKEL, NAIHUAN JING, AND WEIQIANG WANG

Abstract. We establish a q-analog of our recent work on vertex representations and the McKay correspondence. For each finite group Γ we construct a Fock space and associated vertex operators in terms of wreath products of $\Gamma \times \mathbb{C}^\times$ and the symmetric groups. An important special case is obtained when Γ is a finite subgroup of SU_2, where our construction yields a group theoretic realization of the representations of the quantum affine and quantum toroidal algebras of ADE type.

1. Introduction

In our previous paper [FJW] (see [W, FJW] for historical remarks and motivations) we have shown that the basic representation of an affine Lie algebra $\widehat{\mathfrak{g}}$ of ADE type can be constructed from a finite subgroup Γ of SU_2 related to the Dynkin diagram of $\widehat{\mathfrak{g}}$ via the McKay correspondence. In particular, we have recovered a well-known construction [FK, Se] of the basic representation of $\widehat{\mathfrak{g}}$ from the root lattice Q of the corresponding finite dimensional Lie algebra $\mathfrak{g}$. In fact our construction yields naturally the vertex representation of the toroidal Lie algebra $\widehat{\widehat{\mathfrak{g}}}$ which contains the affine Lie algebra as a distinguished subalgebra.

The main goal of the present paper is to q-deform our construction in [FJW]. Again as in the undeformed case we will naturally obtain the earlier construction [FJ] of the basic representation of the quantum affine algebra $U_q(\widehat{\mathfrak{g}})$ from the root lattice Q and its generalization to the quantum toroidal algebra $U_q(\widehat{\widehat{\mathfrak{g}}})$ [GKV] (also cf. [Sa, J3]). The q-deformation is achieved by replacing consistently the representation

---

1991 *Mathematics Subject Classification.* Primary: 17B, 20.

Key words and phrases. finite groups, wreath products, quantum affine algebras and toroidal algebras.

I.F. is supported in part by NSF grant DMS-9700765. N.J. is supported in part by NSA grant MDA904-97-1-0062 and NSF grant DMS-9970493.

theory of Γ by that of $\Gamma \times \mathbb{C}^\times$. The representation ring for $\mathbb{C}^\times$ is identified with the ring of Laurent polynomials $\mathbb{C}[q, q^{-1}]$ so that the formal variable q corresponds to the natural one-dimensional representation of $\mathbb{C}^\times$. It turns out that rather complicated expressions for operators in Drinfeld realization of the quantum affine algebra $U_q(\widehat{\mathfrak{g}})$ and the quantum toroidal algebra $U_q(\widehat{\widehat{\mathfrak{g}}})$ follow instantly from the simple extra factor $\mathbb{C}^\times$. The idea to use representations of $\mathbb{C}^\times$ to obtain a q-deformation of the basic representation was mentioned in [Gr] and is widely used in geometric constructions of representations (see e.g. [CG]).

As in the previous paper [FJW] we give the construction of quantum vertex operators starting from an arbitrary finite group Γ and a self-dual virtual character ξ of $\Gamma \times \mathbb{C}^\times$. Using the restriction and induction functors in representation theory of wreath products of $\Gamma \times \mathbb{C}^\times$ with the symmetric group S_n for all n we construct two "halves" of quantum vertex operators corresponding to any irreducible character γ of $\Gamma \times \mathbb{C}^\times$. Then choosing an irreducible character of $\mathbb{C}^\times$, i.e. an integer power of q we assemble both halves into one quantum vertex operator.

The special case when Γ is a subgroup of SU_2 is important for the application to representation theory of $U_q(\widehat{\mathfrak{g}})$ and for relations [W] to the theory of Hilbert schemes of points on surfaces. To recover the basic representation of $U_q(\widehat{\mathfrak{g}})$ we choose

$$\xi = \gamma_0 \otimes (q + q^{-1}) - \pi \otimes 1_{\mathbb{C}^\times},$$

where γ_0 and $1_{\mathbb{C}^\times}$ are the trivial characters of Γ and $\mathbb{C}^\times$ respectively, q and q^{-1} are the natural and its dual characters of $\mathbb{C}^\times$, and π is the natural character of Γ in SU_2. The fact that the quantum toroidal algebra intrinsically presents in our construction is an additional indication of its importance in representation theory of quantum affine algebras. Moreover when Γ is cyclic of order $r+1$, $\pi \simeq \gamma \oplus \gamma^{-1}$, where γ is the natural character of Γ, one can modify our virtual character ξ with an extra parameter $p = q^k, k \in \mathbb{Z}$ by letting

$$\xi = \gamma_0 \otimes (q + q^{-1}) - (\gamma \otimes p + \gamma^{-1} \otimes p^{-1}).$$

In the special case when $p = q^{\pm 1}$ the quantum vertex representation of the quantum toroidal algebra $U_q(\widehat{\widehat{\mathfrak{g}}})$ can be factored to the basic representation of the quantum affine algebra $U_q(\widehat{\mathfrak{g}})$. This is a q-analog of the factorization in the undeformed case, which exists for an arbitrary simply-laced affine Lie algebra.

To obtain the basic representations of quantum toroidal and affine algebras we only need the quantum vertex operators corresponding to irreducible representations of Γ and their negatives in the Grothendieck ring of this group. We attach two halves of quantum vertex operators using the simplest nontrivial representations of $\mathbb{C}^\times$ namely q and q^{-1}. Each of the two choices and only these two yield the basic representations of $U_q(\widehat{\widehat{\mathfrak{g}}})$ and $U_q(\widehat{\mathfrak{g}})$, in a perfect correspondence with the construction in [FJ]. This choice of an irreducible character of $\mathbb{C}^\times$ is essentially the only freedom that exists in our construction of quantum vertex operators for the quantum affine and toroidal algebras and is fixed by comparison with the algebra relations. However it raises the question of constructing a "natural" quantum vertex operator corresponding to any virtual character γ of Γ. This question is closely related to the well-known problem of finding a q-deformation of vertex operator algebras associated to the basic representation of an affine Lie algebra.

This paper is organized in a way similar to [FJW]. In Sect. 2 we review the theory of wreath products of Γ and extend it to $\Gamma \times \mathbb{C}^\times$. In Sect. 3 we define the weighted bilinear form on $\Gamma \times \mathbb{C}^\times$ and its wreath products. In Sect. 4 we introduce two distinguished q-deformed weight functions associated to subgroups of SU_2. In Sect. 5 we define the Heisenberg algebra associated to Γ and the weighted bilinear form, and we construct its representation in a Fock space. In Sect. 6 we establish the isometry between the representation ring of wreath products of $\Gamma \times \mathbb{C}^\times$ and the Fock space representation of the Heisenberg algebra. In Sect. 7 we construct quantum vertex operators acting on the representation ring of the wreath products. In Sect. 8 we obtain the basic representations of quantum toroidal algebras and quantum affine algebras from representation theory of wreath products for $\Gamma \times \mathbb{C}^\times$.

2. Wreath products and vertex representations

2.1. **The wreath product Γ_n.** Let Γ be a finite group and n a nonnegative integer. The wreath product Γ_n is the semidirect product of the n-th direct product $\Gamma^n = \Gamma \times \cdots \times \Gamma$ and the symmetric group S_n:

$$\Gamma_n = \{(g, \sigma) | g = (g_1, \ldots , g_n) \in \Gamma^n, \sigma \in S_n\}$$

with the group multiplication

$$(g, \sigma) \cdot (h, \tau) = (g\, \sigma(h), \sigma\tau),$$

where S_n acts on Γ^n by permuting the factors.

Let Γ_* be the set of conjugacy classes of Γ consisting of $c^0 = \{1\}$, c^1, $\ldots$, c^r and Γ^* be the set of $r + 1$ irreducible characters: $\gamma_0, \gamma_1, \ldots , \gamma_r$. Here we denote the trivial character of Γ by γ_0. The order of the centralizer of an element in the conjugacy class c is denoted by ζ_c, so the order of the conjugacy class c is $|c| = |\Gamma|/\zeta_c$, where $|\Gamma|$ is the order of Γ.

A partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l)$ is a decomposition of $n = |\lambda| = \lambda_1 + \cdots + \lambda_l$ with nonnegative integers: $\lambda_1 \geq \cdots \geq \lambda_l \geq 1$, where $l = l(\lambda)$ is called the *length* of the partition λ and λ_i are called the *parts* of λ. Another notation for λ is

$$\lambda = (1^{m_1} 2^{m_2} \cdots)$$

with m_i being the multiplicity of parts equal to i in λ. Denote by $\mathcal{P}$ the set of all partitions of integers and by $\mathcal{P}(S)$ the set of all partition-valued functions on a set S. The weight of a partition-valued function $\rho = (\rho(s))_{s \in S}$ is defined to be $\|\rho\| = \sum_{s \in S} |\rho(s)|$. We also denote by $\mathcal{P}_n$ (resp. $\mathcal{P}_n(S)$) the subset of $\mathcal{P}$ (resp. $\mathcal{P}(S)$) of partitions with weight n.

Just as the conjugacy classes of S_n are parameterized by partitions, the conjugacy classes of Γ_n are parameterized by partition-valued functions on Γ_*. Let $x = (g, \sigma) \in \Gamma_n$, where $g = (g_1, \dots, g_n) \in \Gamma^n$ and $\sigma \in S_n$ is presented as a product of disjoint cycles. For each cycle $(i_1 i_2 \cdots i_k)$ of σ, we define the *cycle-product* element $g_{i_k} g_{i_{k-1}} \cdots g_{i_1} \in \Gamma$, which is determined up to conjugacy in Γ by g and the cycle. For any conjugacy class $c \in \Gamma$ and each integer $i \geq 1$, the number of i-cycles in σ whose cycle-product lies in c will be denoted by $m_i(c)$. This gives rise to a partition $\rho(c) = (1^{m_1(c)} 2^{m_2(c)} \dots)$ for $c \in \Gamma_*$. Thus we obtain a partition-valued function $\rho = (\rho(c))_{c \in \Gamma_*} \in \mathcal{P}(\Gamma_*)$ such that $\|\rho\| = \sum_{i,c} i m_i(\rho(c)) = n$. This is called the *type* of the element (g, σ). It is known [M] that two elements in the same conjugacy class have the same type and there exists a one-to-one correspondence between the sets $(\Gamma_n)_*$ and $\mathcal{P}_n(\Gamma_*)$. We will freely say that ρ is the type of the conjugacy class of Γ_n.

Given a class c we denote by c^{-1} the class $\{x^{-1} | x \in c\}$. For each $\rho \in \mathcal{P}(\Gamma_*)$ we also associate the partition-valued function

$$\overline{\rho} = (\rho(c^{-1}))_{c \in \Gamma_*}.$$

Given a partition $\lambda = (1^{m_1} 2^{m_2} \dots)$, we denote by

$$z_\lambda = \prod_{i \geq 1} i^{m_i} m_i!$$

the order of the centralizer of an element of cycle type λ in $S_{|\lambda|}$. The order of the centralizer of an element $x = (g, \sigma) \in \Gamma_n$ of type $\rho = (\rho(c))_{c \in \Gamma_*}$ is given by

$$Z_\rho = \prod_{c \in \Gamma_*} z_{\rho(c)} \zeta_c^{l(\rho(c))}.$$

2.2. **Grothendieck ring** $R_{\Gamma\times C^\times}$**.** Let $R_{\mathbb{Z}}(\Gamma)$ be the $\mathbb{Z}$-lattice generated by γ_i, $i=0,\dots,r$, and $R(\Gamma)=\mathbb{C}\otimes R_{\mathbb{Z}}(\Gamma)$ be the space of complex class functions on the group Γ. In our previous work on the McKay correspondence and vertex representations [W, FJW] we studied the Grothendieck ring $R_\Gamma=\bigoplus_{n\geq 0}R(\Gamma_n)$. In the quantum case we need to add the ring $R(\mathbb{C}^\times)$, the space of characters of $\mathbb{C}^\times=\{t\in\mathbb{C}|t\neq 0\}$.

Let q be the irreducible character of $\mathbb{C}^\times$ that sends t to itself. Then $R(\mathbb{C}^\times)$ is spanned by irreducible multiplicative characters q^n, $n\in\mathbb{Z}$, where

$$q^n(t)=t^n, \qquad t\in\mathbb{C}^\times.$$

Thus $R(\mathbb{C}^\times)$ is identified with the ring $\mathbb{C}[q,q^{-1}]$, and we have

$$R(\Gamma\times\mathbb{C}^\times)=R(\Gamma)\otimes R(\mathbb{C}^\times).$$

An elements of $R(\Gamma\times\mathbb{C}^\times)$ can be written as a finite sum:

$$f=\sum_i f_i\otimes q^{n_i}, \qquad f_i\in R(\Gamma), n_i\in\mathbb{Z}.$$

We can also view f as a function on Γ with values in the ring of Laurent polynomials $\mathbb{C}[q,q^{-1}]$. In this case we will write f^q to indicate the formal variable q, then $f^q(c)=\sum_i f_i(c)q^{n_i}\in\mathbb{C}[q,q^{-1}]$. As a function on $\Gamma\times\mathbb{C}^\times$, we have $f(c,t)=\sum_i f_i(c)t^{n_i}$.

Denote by $R_{\Gamma\times\mathbb{C}^\times}$ the following direct sum:

$$R_{\Gamma\times\mathbb{C}^\times}=\bigoplus_{n\geq 0}R(\Gamma_n\times\mathbb{C}^\times)\simeq R_\Gamma\otimes\mathbb{C}[q,q^{-1}].$$

2.3. **Hopf algebra structure on** $R_{\Gamma\times\mathbb{C}^\times}$**.** The multiplication m in $\mathbb{C}^\times$ and the diagonal map $\mathbb{C}^\times\xrightarrow{d}\mathbb{C}^\times\times\mathbb{C}^\times$ induce the Hopf algebra structure on $R(\mathbb{C}^\times)$.

$$m_{\mathbb{C}^\times}: R(\mathbb{C}^\times)\otimes R(\mathbb{C}^\times)\xrightarrow{\cong}R(\mathbb{C}^\times\times\mathbb{C}^\times)\xrightarrow{d^*}R(\mathbb{C}^\times), \tag{2.1}$$

$$\Delta_{\mathbb{C}^\times}: R(\mathbb{C}^\times)\xrightarrow{m^*}R(\mathbb{C}^\times\times\mathbb{C}^\times)\xrightarrow{\cong}R(\mathbb{C}^\times)\otimes R(\mathbb{C}^\times). \tag{2.2}$$

In terms of the basis $\{q^n\}$ we have

$$q^i\cdot q^j=q^{i+j},$$
$$\Delta(q^k)=q^k\otimes q^k,$$

where we abbreviate $\Delta_{\mathbb{C}^\times}$ by Δ and follow the convention of writing $a\cdot b=m_{\mathbb{C}^\times}(a\otimes b)$.

The antipode $S_{\mathbb{C}^\times}$ and the counit $\epsilon_{\mathbb{C}^\times}$ are given by

$$S_{\mathbb{C}^\times}(q^n)=q^{-n}, \qquad \epsilon_{\mathbb{C}^\times}(q^n)=\delta_{n0}.$$

We extend the Hopf algebra structures on $R(\mathbb{C}^\times)$ and R_Γ [Z, M] into a Hopf algebra structure on $R_{\Gamma\times\mathbb{C}^\times}$ using a standard procedure in Hopf algebra [A]. The multiplication and comultiplication are given by the respective composition of the following maps:

$$\begin{aligned} m : R(\Gamma_n \times \mathbb{C}^\times) \otimes R(\Gamma_m \times \mathbb{C}^\times) &\xrightarrow{\cong} R(\Gamma_n \times \mathbb{C}^\times \times \Gamma_m \times \mathbb{C}^\times) \\ &\xrightarrow{1\otimes m_{\mathbb{C}^\times}} R(\Gamma_n \times \Gamma_m \times \mathbb{C}^\times) \xrightarrow{Ind\otimes 1} R(\Gamma_{n+m} \times \mathbb{C}^\times); \qquad (2.3) \\ \Delta : R(\Gamma_n \times \mathbb{C}^\times) &\xrightarrow{Res\otimes 1} \oplus_{m=0}^n R(\Gamma_{n-m} \times \Gamma_m \times \mathbb{C}^\times) \\ &\xrightarrow{1\otimes \Delta_{\mathbb{C}^\times}} \oplus_{m=0}^n R(\Gamma_{n-m} \times \Gamma_m \times \mathbb{C}^\times \times \mathbb{C}^\times) \\ &\xrightarrow{\cong} \oplus_{m=0}^n R(\Gamma_{n-m} \times \mathbb{C}^\times) \otimes R(\Gamma_m \times \mathbb{C}^\times), \qquad (2.4) \end{aligned}$$

where we have used the identification of $R(\mathbb{C}^\times \times \mathbb{C}^\times)$ with $R(\mathbb{C}^\times) \otimes R(\mathbb{C}^\times)$ in (2.1-2.2). Also $Ind : R(\Gamma_n \times \Gamma_m) \longrightarrow R(\Gamma_{n+m})$ denotes the induction functor and $Res : R(\Gamma_n) \longrightarrow R(\Gamma_{n-m} \times \Gamma_m)$ denotes the restriction functor.

The antipode is given by

$$S(f(g,t)) = f(g^{-1}, t^{-1}), \qquad g \in \Gamma, t \in \mathbb{C}^\times.$$

In particular, $S(\gamma)(c) = \gamma(c^{-1})$ for $\gamma \in \Gamma^*$. As we mentioned earlier, we may write $f \in R_{\Gamma\times\mathbb{C}^\times}$ as

$$f^q(g) = \sum_i f_i(g) q^{n_i},$$

Then $S(f^q)(g) = \sum_i f_i(g^{-1}) q^{-n_i}$.

The counit ϵ is defined by

$$\epsilon(R(\Gamma_n \times \mathbb{C}^\times)) = 0, \qquad \text{if} \quad n \neq 0,$$

and ϵ on $R(\mathbb{C}^\times)$ is the counit of the Hopf algebra $R(\mathbb{C}^\times)$.

3. A WEIGHTED BILINEAR FORM ON $R(\Gamma_n \times \mathbb{C}^\times)$

3.1. **A standard bilinear form on** $R_{\Gamma\times\mathbb{C}^\times}$**.** Let $f, g \in R(\Gamma\times\mathbb{C}^\times)$ with $f = \sum_i f_i \otimes q^{n_i}$ and $g = \sum_i g_i \otimes q^{m_i}$. The $\mathbb{C}[q, q^{-1}]$-valued standard $\mathbb{C}$-bilinear form on $R(\Gamma \times \mathbb{C}^\times)$ is defined as

$$\begin{aligned} \langle f, g\rangle_\Gamma^q &= \sum_{i,j} \langle f_i, g_j\rangle_\Gamma q^{n_i - m_j} \\ &= \sum_{i,j} \sum_{c\in\Gamma_*} \zeta_c^{-1} f_i(c) g_j(c^{-1}) q^{n_i - m_j}, \end{aligned}$$

where we recall that c^{-1} denotes the conjugacy class $\{x^{-1}|x \in c\}$ of Γ, and ζ_c is the order of the centralizer of the class c in Γ. Sometimes we will also view the bilinear form as a function of $t \in \mathbb{C}^\times$:

$$\langle f, g\rangle_\Gamma^q(t) = \sum_{c\in\Gamma_*} \zeta_c^{-1} f(c,t) S(g(c,t)).$$

The following is a direct consequence of the orthogonality of irreducible characters of Γ.

$$\begin{aligned} \langle \gamma_i \otimes q^k, \gamma_j \otimes q^l\rangle_\Gamma^q &= \delta_{ij} q^{k-l}, \\ \sum_{\gamma\in\Gamma^*} \gamma(c')S(\gamma)(c) &= \delta_{c,c'}\zeta_c, \quad c, c' \in \Gamma_*. \end{aligned} \tag{3.1}$$

Let $\langle\ ,\ \rangle_{\Gamma_n}^q$ be the $\mathbb{C}[q,q^{-1}]$-valued bilinear form on $R(\Gamma_n \times \mathbb{C}^\times)$. The $\mathbb{C}[q,q^{-1}]$-valued standard bilinear form in $R_{\Gamma\times\mathbb{C}^\times}$ is defined in terms of the bilinear form on $R(\Gamma_n \times \mathbb{C}^\times)$ as follows:

$$\langle u, v\rangle^q = \sum_{n\geq 0} \langle u_n, v_n\rangle_{\Gamma_n}^q,$$

where $u = \sum_n u_n$ and $v = \sum_n v_n$ with $u_n, v_n \in R(\Gamma_n \times \mathbb{C}^\times)$.

3.2. **A weighted bilinear form on $R(\Gamma \times \mathbb{C}^\times)$.** A class function $\xi \in R(\Gamma \times \mathbb{C}^\times)$ is called self-dual if for all $x \in \Gamma, t \in \mathbb{C}^\times$

$$\xi(x,t) = S(\xi(x,t)),$$

or equivalently $\xi^q(x) = \xi^{q^{-1}}(x^{-1})$.

We fix a self-dual class function ξ. The tensor product of two representations γ and β in $R(\Gamma \times \mathbb{C}^\times)$ will be denoted by $\gamma * \beta$.

Let $a_{ij} \in \mathbb{C}[q,q^{-1}]$ be the (virtual) multiplicity of γ_j in $\xi * \gamma_i$, i.e.,

$$\xi * \gamma_i = \sum_{j=0}^r a_{ij}\gamma_j. \tag{3.2}$$

We denote by A^q the $(r+1)\times(r+1)$ matrix $(a_{ij})_{0\leq i,j\leq r}$.

Associated to ξ we introduce the following weighted bilinear form

$$\langle f, g\rangle_\xi^q = \langle \xi * f, g\rangle_\Gamma^q, \quad f, g \in R(\Gamma \times \mathbb{C}^\times).$$

where we use the superscript q to indicate the q-dependence. The superscript q is often omitted if the q-variable in characters f and g is clear from the context. The explicit formula of the bilinear form is given as follows.

$$\begin{aligned}\langle f, g\rangle_\xi^q &= \frac{1}{|\Gamma|}\sum_{x\in\Gamma}\xi^q(x)f^q(x)g^{q^{-1}}(x^{-1})\\ &= \sum_{c\in\Gamma_*}\zeta_c^{-1}\xi^q(c)f^q(c)g^{q^{-1}}(c^{-1}),\end{aligned}\tag{3.3}$$

which is the average of the character $\xi * f * \overline{g}$ over Γ.

The self-duality of ξ together with (3.3) implies that

$$a_{ij} = \overline{a_{ji}},$$

i.e. A^q is a hermitian-like matrix with the bar action given by $\overline{q} = q^{-1}$.

The orthogonality (3.1) implies that

$$a_{ij} = \langle\gamma_i, \gamma_j\rangle_\xi^q.\tag{3.4}$$

Remark 3.1. If ξ is the trivial character γ_0, then the weighted bilinear form becomes the standard one on $R(\Gamma \times \mathbb{C}^\times)$.

3.3. **A weighted bilinear form on $R(\Gamma_n \times \mathbb{C}^\times)$.** Let V be a $\Gamma\times\mathbb{C}^\times$-module which affords a character γ in $R(\Gamma \times \mathbb{C}^\times)$. We can decompose V as follows:

$$V = \bigoplus_i V_i \otimes \mathbb{C}(k_i),$$

where V_i is a (virtual) Γ-module in $R(\Gamma)$ and $\mathbb{C}(k_i)$ is the one dimensional $\mathbb{C}^\times$-module afforded by the character q^{k_i}.

The n-th outer tensor product $V^{\otimes n}$ of V can be regarded naturally as a representation of the wreath product $(\Gamma \times \mathbb{C}^\times)_n$ via permutation of the factors and the usual direct product action. More precisely, note that $\Gamma_n \times \mathbb{C}^\times$ can be viewed as a subgroup of $(\Gamma\times\mathbb{C}^\times)_n$ by the diagonal inclusion from $\mathbb{C}^\times$ to $(\mathbb{C}^\times)^n$:

$$\Gamma_n \times \mathbb{C}^\times \longrightarrow (\Gamma^n \times {\mathbb{C}^\times}^n) \rtimes S_n = (\Gamma \times \mathbb{C}^\times)_n.$$

This provides a natural $\Gamma_n \times \mathbb{C}^\times$-module structure on $V^{\otimes n}$. We denote its character by $\eta_n(\gamma)$. Explicitly we have

$$(g, \sigma, t).(v_1 \otimes \cdots \otimes v_n) = (g_1, t)v_{\sigma^{-1}(1)} \otimes \cdots \otimes (g_n, t)v_{\sigma^{-1}(n)},\tag{3.5}$$

where $g = (g_1, \dots, g_n) \in \Gamma^n$.

Let ε_n be the (1-dimensional) sign representation of Γ_n so that Γ^n acts trivially while letting S_n act as a sign representation. We denote by $\varepsilon_n(\gamma) \in R(\Gamma_n \times \mathbb{C}^\times)$ the character of the tensor product of $\varepsilon_n \otimes 1$ and $V^{\otimes n}$.

The weighted bilinear form on $R(\Gamma_n \times \mathbb{C}^\times)$ is now defined by

$$\langle f, g\rangle_{\xi,\Gamma_n}^q = \langle\eta_n(\xi) * f, g\rangle_{\Gamma_n}^q, \quad f, g \in R(\Gamma_n \times \mathbb{C}^\times).$$

We shall see in Corollary 6.2 that $\eta_n(\xi)$ is self-dual if the class function ξ is invariant under the antipode S. In such a case the matrix of the bilinear form $\langle\ ,\ \rangle_\xi^q$ is equal to its adjoint (transpose and bar action).

We can naturally extend η_n to a map from $R(\Gamma)\otimes q^k$ to $R(\Gamma\times\mathbb{C}^\times)$ as in the classical case (cf. [W]). In particular, if β and γ are characters of representations V and W of Γ respectively, then

$$\begin{aligned}&\eta_n(\beta\otimes q^k+\gamma\otimes q^l)\\ &=\sum_{m=0}^{n} Ind_{\Gamma_{n-m}\times\mathbb{C}^\times\times\Gamma_m\times\mathbb{C}^\times}^{\Gamma_n\times\mathbb{C}^\times}[\eta_{n-m}(\beta\otimes q^k)\otimes\eta_m(\gamma\otimes q^l)],\end{aligned}\tag{3.6}$$

$$\begin{aligned}&\eta_n(\beta\otimes q^k-\gamma\otimes q^l)\\ &=\sum_{m=0}^{n}(-1)^m Ind_{\Gamma_{n-m}\times\mathbb{C}^\times\times\Gamma_m\times\mathbb{C}^\times}^{\Gamma_n\times\mathbb{C}^\times}[\eta_{n-m}(\beta\otimes q^k)\otimes\varepsilon_m(\gamma\otimes q^l)].\end{aligned}\tag{3.7}$$

On $R_{\Gamma\times\mathbb{C}^\times}=\bigoplus_n R(\Gamma_n\times\mathbb{C}^\times)$ the weighted bilinear form is given by

$$\langle u,v\rangle_\xi^q=\sum_{n\geq 0}\langle u_n,v_n\rangle_{\xi,\Gamma_n}^q$$

where $u=\sum_n u_n$ and $v=\sum_n v_n$ with $u_n,v_n\in R(\Gamma_n\times\mathbb{C}^\times)$.

The bilinear form $\langle\ ,\ \rangle_\xi^q$ on $R_{\Gamma\times\mathbb{C}^\times}$ is $\mathbb{C}$-bilinear and takes values in $\mathbb{C}[q,q^{-1}]$. When $n=1$, it reduces to the weighted bilinear form defined on $R(\Gamma\times\mathbb{C}^\times)$.

We will often omit the superscript q and use the notation $\langle\ ,\ \rangle_\xi$ for the weighted bilinear form on $R_{\Gamma\times\mathbb{C}^\times}$.

4. Quantum McKay weights

4.1. **Quantum McKay correspondence.** Let $d_i=\gamma_i(c^0)$ be the dimension of the irreducible representation of Γ corresponding to the character γ_i.

The following generalizes a result of McKay [Mc].

Proposition 4.1. *For each class $c\in\Gamma_*$ the column vector*

$$v(c)=(\gamma_0(c),\gamma_1(c),\dots,\gamma_r(c))^t$$

is an eigenvector of the $(r+1)\times(r+1)$-matrix $A^q=(\langle\gamma_i,\gamma_j\rangle_\xi^q)$ with eigenvalue $\xi^q(c)$. In particular $(d_0,d_1,\dots,d_r)$ is an eigenvector of A^q with eigenvalue $\xi^q(c^0)$.

Proof. We compute directly that

$$\begin{aligned}\sum_{k=0}^{r}\langle\gamma_i,\gamma_k\rangle_\xi^q\gamma_k(c) &= \sum_k\sum_{c'\in\Gamma_*}\zeta_{c'}^{-1}\xi^q(c')\gamma_i(c')\gamma_k(c'^{-1})\gamma_k(c)\\ &= \sum_{c'\in\Gamma_*}\zeta_{c'}^{-1}\xi^q(c')\gamma_i(c')\sum_k\gamma_k(c'^{-1})\gamma_k(c)\\ &= \sum_{c'\in\Gamma_*}\zeta_{c'}^{-1}\xi^q(c')\gamma_i(c')\zeta_c\delta_{cc'}\\ &= \xi^q(c)\gamma_i(c).\end{aligned}$$

□

Let π be an irreducible faithful representation π of Γ of dimension d. For each integer n we define the q-integer $[n]$ that can be viewed as a character of $\mathbb{C}^\times$ by

$$[n]=\frac{q^n-q^{-n}}{q-q^{-1}}=q^{n-1}+q^{n-3}+\cdots+q^{-n+1}.$$

We take the following special class function

$$\xi=\gamma_0\otimes[d]-\pi\otimes 1_{\mathbb{C}^\times}, \tag{4.1}$$

where we have also used the symbol π for the corresponding character, and $1_{\mathbb{C}^\times}=q^0$ is the trivial character of $\mathbb{C}^\times$.

Proposition 4.2. *The weighted bilinear form associated to (4.1) is non-degenerate. If π is an embedding of Γ into SU_d and $t\neq 1$ is a nonnegative real number, then the weighted bilinear form evaluated on t is positive definite.*

Proof. A simple fact of finite group theory says that

$$\langle f,f\rangle_\pi\le d\langle f,f\rangle.$$

Assume that $t\in\mathbb{R}_+$. Observe that

$$t^{d-1}+t^{d-3}+\cdots+t^{-d+1}\ge d$$

and the equality holds if and only if $t=1$.

Let $A^q=(\langle\gamma_i,\gamma_j\rangle_\xi^q)=(a_{ij})$. Note that for any faithful representation π of Γ we have that

$$\pi*\gamma_i=\sum_j c_{ij}\gamma_j,\qquad c_{ij}\in\mathbb{N}.$$

Then it follows that

$$\begin{aligned}\langle\gamma_i,\gamma_j\rangle_\xi^q(t) &= (t^{d-1}+t^{d-3}+\cdots+t^{-d+1})\langle\gamma_i,\gamma_j\rangle-\langle\gamma_i,\gamma_j\rangle_\pi\\ &= [d](t)\delta_{ij}-c_{ij}=A^1+([d](t)-d)I.\end{aligned}$$

According to Steinberg (see e.g. [FJW]), A^1 is positive semi-definite which generalizes McKay's observation in the case of $d = 2$. This implies that the eigenvalues of A^q are $\geq [d](t) - d \geq 0$. Thus the matrix $A^q(t)$ is positive-definite when $t > 0$ and $t \neq 1$. $\square$

We remark that when $|t| = 1$ and t is close to 1, the signature of $A^q(t)$ is $(-1, 1, \ldots, 1)$ due to $[d](t) \leq d$.

Remark 4.1. The matrix A^1 is integral, and the entries of A^q are the q-numbers of the corresponding entries in A^1 when $r \geq 2$.

4.2. **Two quantum McKay weights.** Let Γ is a finite subgroup of SU_2 and we introduce the first distinguished self-dual class function

$$\xi = \gamma_0 \otimes (q + q^{-1}) - \pi \otimes 1_{\mathbb{C}^\times},$$

where π is the character of the embedding of Γ in SU_2.

The matrix of the weighted bilinear form $\langle\ ,\ \rangle_\xi$ (cf. (3.4)) has the following entries:

$$a_{ij} = \begin{cases} q + q^{-1}, & \text{if } i = j, \\ -1, & \text{if } \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1, \\ -2, & \text{if } \langle \gamma_i, \gamma_j \rangle_\xi^1 = -2 \text{ and } \Gamma = \mathbb{Z}/2\mathbb{Z} \\ 0, & \text{otherwise.} \end{cases} \tag{4.2}$$

In particular when $q = 1$ the matrix (a_{ij}^1) coincides with the extended Cartan matrix of ADE type according to the five classes of finite subgroups of SU_2: the cyclic, binary dihedral, tetrahedral, octahedral, and icosahedral groups. McKay [Mc] gave a direct correspondence between a finite subgroup of SU_2 and the affine Dynkin diagram D of ADE type. Each irreducible character γ_i corresponds to a vertex of D, and the number of edges between γ_i and γ_j $(i \neq j)$ is equal to $|\langle \gamma_i, \gamma_j \rangle_\xi^1|$, where $\langle \gamma_i, \gamma_j \rangle_\xi^1 = a_{ij}^1$ are the entries of matrix A^1 of the weighted bilinear form $\langle\ ,\ \rangle_\xi^1$. For this reason we will call our matrix $A^q = (a_{ij}) = (\langle \gamma_i, \gamma_j \rangle_\xi^q)$ the quantum Cartan matrix.

Let Γ be the cyclic subgroup of SU_2 of order $r+1$. We can introduce the second deformation parameter in the quantum Cartan matrix. Let $\gamma_i (i = 0, \ldots, r)$ be the full set of irreducible characters of $\mathbb{C}^\times$ such that $\gamma_i * \gamma_j = \gamma_{i+j \mod r+1}$. The embedding of Γ in SU_2 is given by $\pi = \gamma_1 + \gamma_r$.

For $p = q^k \in R(\mathbb{C}^\times)$ we let

$$\xi = \xi^{q,p} = \gamma_0 \otimes (q + q^{-1}) - (\gamma_1 \otimes p + \gamma_r \otimes p^{-1}).$$

When $p = 1$ the second choice reduces to the first choice in type A. This class function is self-dual since $S(\gamma_i) = \gamma_{r+1-i}, i = 0, 1, \ldots, r$ and $S(q) = q^{-1}, S(p) = p^{-1}$.

It is easy to see that

$$a_{ij}(q,p)=\langle\gamma_i,\gamma_j\rangle_\xi^{q,p}=[2]\delta_{ij}-p\delta_{i+1,j}-p^{-1}\delta_{i-1,j}. \tag{4.3}$$

Thus the matrix of the weighted bilinear form $\langle\ ,\ \rangle_\xi$ (cf. (3.4)) has the following form.

$$\begin{pmatrix} q+q^{-1} & -p & 0 & \cdots & -p^{-1} \\ -p^{-1} & q+q^{-1} & -p & \cdots & 0 \\ 0 & -p^{-1} & q+q^{-1} & \cdots & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ -p & 0 & \cdots & -p^{-1} & q+q^{-1} \end{pmatrix}, \qquad \text{if } r\geq 2, \tag{4.4}$$

or

$$\begin{pmatrix} q+q^{-1} & -p-p^{-1} \\ -p-p^{-1} & q+q^{-1} \end{pmatrix}, \qquad \text{if } r=1. \tag{4.5}$$

Note that when $\Gamma=1$, the matrix of the bilinear form $\langle\ ,\ \rangle_\xi^{q,p}$ is $q+q^{-1}-p-p^{-1}$, which is degenerate when $q=p^{\pm1}$.

We will call this matrix the (q,p)-Cartan matrix (of type A). The self-duality of $\xi^{q,p}$ transforms into the condition that the (q,p)-Cartan matrix is $*$-invariant, where the $*$ action is the composition of transpose and bar action. Namely, $a_{ij}(q,p)=a_{ji}(q^{-1},p^{-1})$.

Proposition 4.3. *If $p\neq q^{\pm1}$, then the bilinear form $\langle\ ,\ \rangle_\xi^{q,p}$ is non-degenerate. If $p=q^{\pm1}$, the bilinear form $\langle\ ,\ \rangle_\xi^{q,p}$ is degenerate of rank r.*

Proof. Let $A^{q,p}=(\langle\gamma_i,\gamma_j\rangle_\xi^{q,p})$ be the matrix of the bilinear form $\langle\ ,\ \rangle_\xi^{q,p}$ and let ω be a $(r+1)$-th root of unity. Then $\gamma_i(c^j)=\omega^{ij}$ and $\gamma_i*\gamma_j=\gamma_{i+j}$. From this and Proposition 4.1 we see that as a matrix over $\mathbb{C}[q,q^{-1}]$ the eigenvalues of $A^{p,q}$ are $q+q^{-1}-\omega^ip-\omega^{-i}p^{-1}$, $i=0,\dots,r$. The function $q+q^{-1}-\omega^ip-\omega^{-i}p^{-1}\in R(\mathbb{C}^\times)$ is non-zero except when $i=0$ and $q=p^{\pm1}$. □

5. Quantum Heisenberg algebras and Γ_n

5.1. **Heisenberg algebra $\widehat{\mathfrak{h}}_{\Gamma,\xi}$.** Let $\widehat{\mathfrak{h}}_{\Gamma,\xi}$ be the infinite dimensional Heisenberg algebra over $\mathbb{C}[q,q^{-1}]$, associated with Γ and $\xi\in R(\Gamma\times\mathbb{C}^\times)$, with generators $a_m(c), c\in\Gamma_*, m\in\mathbb{Z}$ and a central element C subject to the following commutation relations:

$$[a_m(c^{-1}),a_n(c')]=m\delta_{m,-n}\delta_{c,c'}\zeta_c\xi_{q^m}(c)C, \quad c,c'\in\Gamma_*. \tag{5.1}$$

For $m\in\mathbb{Z},\gamma\in\Gamma^*$ and $k\in\mathbb{Z}$ we define

$$a_m(\gamma\otimes q^k)=\sum_{c\in\Gamma_*}\zeta_c^{-1}\gamma(c)a_m(c)q^{mk}$$

and then extend it to $R(\Gamma \times \mathbb{C}^\times)$ linearly over $\mathbb{C}$. Thus we have for $\gamma \in R(\Gamma \times \mathbb{C}^\times)$

$$a_m(\gamma) = \sum_{c \in \Gamma_*} \zeta_c^{-1} \gamma_{q^m}(c) a_m(c). \tag{5.2}$$

In particular we have $a_m(\gamma \otimes q^k) = a_m(\gamma) q^{mk}$.

It follows immediately from the orthogonality (3.1) of the irreducible characters of Γ that for each $c \in \Gamma_*$

$$a_m(c) = \sum_{\gamma \in \Gamma^*} S(\gamma(c)) a_m(\gamma).$$

Note that this formula is also valid if the summation runs through $\Gamma^* \otimes q^k$ with a fixed k.

Proposition 5.1. *The Heisenberg algebra* $\widehat{\mathfrak{h}}_{\Gamma,\xi}$ *has a new basis given by* $a_n(\gamma)$ *and* C $(n \in \mathbb{Z}, \gamma \in \Gamma^*)$ *over* $\mathbb{C}[q, q^{-1}]$ *with the following relations:*

$$[a_m(\gamma), a_n(\gamma')] = m\delta_{m,-n} \langle \gamma, \gamma' \rangle_\xi^{q^m} C. \tag{5.3}$$

Proof. This is proved by a direct computation using Eqns. (5.1), (3.3) and (3.1).

$$\begin{aligned}
[a_m(\gamma), a_n(\gamma')] &= \sum_{c,c' \in \Gamma_*} \zeta_c^{-1} \zeta_{c'}^{-1} \gamma(c) \gamma'(c') [a_m(c), a_n(c')] \\
&= m\delta_{m,-n} \sum_{c,c' \in \Gamma_*} \zeta_c^{-1} \zeta_{c'}^{-1} \gamma(c) \gamma'(c') \delta_{c^{-1},c'} \zeta_c \xi_{q^m}(c) C \\
&= m\delta_{m,-n} \sum_{c \in \Gamma_*} \zeta_c^{-1} \gamma(c) \gamma'(c^{-1}) \xi_{q^m}(c) C \\
&= m\delta_{m,-n} \langle \gamma, \gamma' \rangle_\xi^{q^m} C.
\end{aligned}$$

□

5.2. **Action of $\widehat{\mathfrak{h}}_{\Gamma,\xi}$ on the Space $S_{\Gamma \times \mathbb{C}^\times}$.** Let $S_{\Gamma \times \mathbb{C}^\times}$ be the symmetric algebra generated by $a_{-n}(\gamma), n \in \mathbb{N}, \gamma \in \Gamma_*$ over $\mathbb{C}[q, q^{-1}]$. We define $a_{-n}(\gamma \otimes q^k) = a_{-n}(\gamma) q^{-kn}$ and the natural degree operator on the space $S_{\Gamma \times \mathbb{C}^\times}$ by

$$\deg(a_{-n}(\gamma \otimes q^k)) = n$$

which makes $S_{\Gamma \times \mathbb{C}^\times}$ into a $\mathbb{Z}_+$-graded algebra.

The space $S_{\Gamma \times \mathbb{C}^\times}$ affords a natural realization of the Heisenberg algebra $\widehat{\mathfrak{h}}_{\Gamma,\xi}$ with $C = 1$. Since $a_{-n}(\gamma \otimes q^k) = q^{-nk} a_{-n}(\gamma)$, it is enough to describe the action for $a_{-n}(\gamma)$. The central element C acts as the identity operator. For $n > 0$, $a_{-n}(\gamma)$ act as multiplication operators on $S_{\Gamma\times\mathbb{C}^\times}$. The element $a_n(\gamma), n \geq 0$ acts as a differential operator through contraction:

$$\begin{aligned} &a_n(\gamma).a_{-n_1}(\alpha_1)a_{-n_2}(\alpha_2)\dots a_{-n_k}(\alpha_k) \\ &= \textstyle\sum_{i=1}^k \langle\gamma,\alpha_i\rangle_\xi^{q^n} a_{-n_1}(\alpha_1)a_{-n_2}(\alpha_2)\dots \check{a}_{-n_i}(\alpha_i)\dots a_{-n_k}(\alpha_k). \end{aligned}$$

Here $n_i > 0, \alpha_i \in R(\Gamma)$ for $i = 1, \dots, k$, and $\check{a}_{-n_i}(\alpha_i)$ means the very term is deleted. In this case $S_{\Gamma\times\mathbb{C}^\times}$ is an irreducible representation of $\widehat{\mathfrak{h}}_{\Gamma,\xi}$ with the unit 1 as the highest weight vector.

5.3. **The bilinear form on $S_{\Gamma\times\mathbb{C}^\times}$.** As a $\widehat{\mathfrak{h}}_{\Gamma,\xi}$-module, the space $S_{\Gamma\times\mathbb{C}^\times}$ admits a bilinear form $\langle\ ,\ \rangle'_\xi$ over $\mathbb{C}[q,q^{-1}]$ characterized by

$$\begin{aligned} &\langle 1,1\rangle'_\xi = 1, \\ &\langle au,v\rangle'_\xi = \langle u,a^*v\rangle'_\xi, \qquad a\in\widehat{\mathfrak{h}}_{\Gamma,\xi}, \end{aligned} \tag{5.4}$$

with the adjoint map $*$ on $\widehat{\mathfrak{h}}_{\Gamma,\xi}$ given by

$$a_n(\gamma\otimes q^k)^* = a_{-n}(\gamma\otimes q^k), \qquad n\in\mathbb{Z}. \tag{5.5}$$

Note that the adjoint map $*$ is a $\mathbb{C}$-linear anti-homomorphism of $\widehat{\mathfrak{h}}_{\Gamma,\xi}$, and $q^* = \overline{q}$. We still use the same symbol $*$ to denote the hermitian-like dual, since it clearly generalizes the $*$-action on the deformed Cartan matrix (4.4).

For any partition $\lambda = (\lambda_1, \lambda_2, \dots)$ and $\gamma\in\Gamma^*$, we define

$$a_{-\lambda}(\gamma) = a_{-\lambda_1}(\gamma)a_{-\lambda_2}(\gamma)\dots.$$

For $\rho = (\rho(\gamma))_{\gamma\in\Gamma^*}\in\mathcal{P}(\Gamma^*)$, we define

$$a_{-\rho\otimes q^k} = q^{-k\|\rho\|}\prod_{\gamma\in\Gamma^*} a_{-\rho(\gamma)}(\gamma).$$

It is clear that for a fixed $k\in\mathbb{Z}$ the elements $a_{-\rho\otimes q^k}, \rho\in\mathcal{P}(\Gamma^*)$ form a basis of $S_{\Gamma\times\mathbb{C}^\times}$ over $\mathbb{C}[q,q^{-1}]$.

Given a partition $\lambda = (\lambda_1, \lambda_2, \dots)$ and $c\in\Gamma_*$, we define

$$a_{-\lambda}(c\otimes q^k) \ = \ q^{-k|\lambda|}a_{-\lambda_1}(c)a_{-\lambda_2}(c)\dots,$$

For any $\rho = (\rho(c))_{c\in\Gamma_*}\in\mathcal{P}(\Gamma_*)$ and $k\in\mathbb{Z}$, we define

$$a'_{-\rho\otimes q^k} = q^{-k\|\rho\|}\prod_{c\in\Gamma_*} a_{-\rho(c)}(c).$$

It follows from Proposition 5.1 that

$$\langle a'_{-\rho\otimes q^k}, a'_{-\overline{\sigma}\otimes q^l}\rangle'_\xi = \delta_{\rho,\sigma}q^{\|\rho\|(l-k)}Z_\rho\prod_{c\in\Gamma_*}\prod_{i\geq 1}\xi_{q^i}(c)^{m_i(\rho(c))}, \tag{5.6}$$

where $\rho, \sigma \in \mathcal{P}(\Gamma_*)$. Note that $S(a'_{-\rho\otimes q^k}) = a'_{-\overline{\rho}\otimes q^{-k}}$, where we recall that $\overline{\rho} \in \mathcal{P}(\Gamma_*)$ is the partition-valued function given by $c \mapsto \rho(c^{-1})$, $c \in \Gamma$.

6. The characteristic map as an isometry

6.1. **The characteristic map ch.** Let $\Psi : \Gamma_n \to S_{\Gamma\times\mathbb{C}^\times}$ be the map defined by $\Psi(x) = a'_{-\rho}$ if $x \in \Gamma_n$ is of type ρ.

We define a $\mathbb{C}$-linear map $ch : R_{\Gamma\times\mathbb{C}^\times} \longrightarrow S_{\Gamma\times\mathbb{C}^\times}$ by letting

$$\begin{aligned} ch(f) &= \langle f, \Psi\rangle_{\Gamma_n} \\ &= \sum_{\rho\in\mathcal{P}(\Gamma_*)} Z_\rho^{-1} S(f(\rho)) a'_{-\rho}, \end{aligned} \tag{6.1}$$

where $f(\rho) \in \mathbb{C}[q, q^{-1}]$ is the value of f at the elements of type ρ. The map ch is called the *characteristic map*. This generalizes the definition of the characteristic map in the classical setting (cf. [M, FJW]).

The space $S_{\Gamma\times\mathbb{C}^\times}$ can also be interpreted as follows. The element $a_{-n}(\gamma), n > 0, \gamma \in \Gamma^*$ is identified as the n-th power sum in a sequence of variables $y_g = (y_{i\gamma})_{i\geq 1}$. By the commutativity among $a_{-n}(\gamma)$ ($\gamma \in \Gamma^*, n > 0$) and dimension counting it is clear that the space $S_{\Gamma\times\mathbb{C}^\times}$ is isomorphic with the space Λ_Γ of symmetric functions indexed by Γ^* tensored with $\mathbb{C}[q, q^{-1}]$ (cf. [M]).

Denote by $c_n (c \in \Gamma_*)$ the conjugacy class in Γ_n of elements $(x, s) \in \Gamma_n$ such that s is an n-cycle and $x \in c$. Denote by $\sigma_n(c \otimes q^k)$ the class function on $\Gamma_n \times \mathbb{C}^\times$ which takes values $n\zeta_c t^{-nk}$ (i.e. the order of the centralizer of an element in the class c_n times t^{-nk}) on elements in the class $c_n \times t$ and 0 elsewhere. For $\rho = \{m_r(c)\}_{r\geq 1, c\in\Gamma_*} \in \mathcal{P}_n(\Gamma^*)$ and $k \in \mathbb{Z}$,

$$\sigma_{\rho\otimes q^k} = q^{-nk} \prod_{r\geq 1, c\in\Gamma_*} \sigma_r(c)^{m_r(c)}$$

is the class function on $\Gamma_n \times \mathbb{C}^\times$ which takes value $Z_\rho t^{-nk}$ on the conjugacy class of type $\rho \times t$ and 0 elsewhere. Given $\gamma \in \Gamma^*$ and $k \in \mathbb{Z}$, we denote by $\sigma_n(\gamma \otimes q^k)$ the class function on $\Gamma_n \times \mathbb{C}^\times$ which takes values $n\gamma(c)t^{-nk}$ on elements in the class $c_n \times t (c \in \Gamma_*)$ and 0 elsewhere.

Lemma 6.1. *The map ch sends $\sigma_{\rho\otimes q^k}$ to $a'_{-\rho\otimes q^k}$. In particular, it sends $\sigma_n(\gamma \otimes q^k)$ to $a_{-n}(\gamma \otimes q^k)$ in $S_{\Gamma\times\mathbb{C}^\times}$.*

Proof. This is verified by the definition of ch (6.1) and the character values of σ_n defined above. □

Proposition 6.1. *Given $\gamma \in \Gamma^*$, the character value of $\eta_n(\gamma \otimes q^k)$ on the conjugacy class c_ρ of type $\rho = (\rho(c))_{c \in \Gamma_*}$ is given by*

$$\eta_n(\gamma \otimes q^k)(c_\rho) = \prod_{c \in \Gamma_*} \gamma(c)^{l(\rho(c))} q^{nk}. \tag{6.2}$$

In particular, we have $\eta_n(\gamma \otimes q^k) = \eta_n(\gamma) q^{nk}$.

Proof. We first let (g, σ) be an element of Γ_n such that σ is a cycle of length n, say $\sigma = (12 \cdots n)$. Let $\{e_i\}$ be a basis of V, and $\gamma \otimes q^k$ is afforded by the action: $(h, t) e_j = \sum_i c_{ij}(h) t^k e_i$, where $h \in \Gamma$. We then have

$$\begin{aligned}
&(g, \sigma, t).(e_{j_1} \otimes e_{j_2} \otimes \cdots \otimes e_{j_n}) \\
&= (g_1, t) e_{j_n} \otimes (g_2, t) e_{j_1} \otimes \cdots \otimes (g_n, t) e_{j_{n-1}} \\
&= \sum_{i_1, \ldots, i_n} t^{kn} c_{i_n j_n}(g_1) c_{i_1 j_1}(g_2) \cdots c_{i_{n-1} j_{n-1}}(g_n) e_{i_n} \otimes e_{i_1} \cdots \otimes e_{i_{n-1}}.
\end{aligned}$$

It follows that

$$\begin{aligned}
\eta_n(\gamma \otimes q^k)(c_\rho, t) &= \text{trace } (g, \sigma, t) \\
&= \sum_{j_1, \ldots, j_n} t^{kn} c_{j_1 j_n}(g_1) c_{j_2 j_1}(g_2) \cdots c_{j_n j_{n-1}}(g_n) \\
&= \text{trace } t^{kn} a(g_n) a(g_{n-1}) \ldots a(g_1) \\
&= \text{trace } t^{kn} a(g_n g_{n-1} \ldots g_1) = \gamma(c) q^{kn}(t).
\end{aligned}$$

Given $x \times y \in \Gamma_n$ where $x \in \Gamma_r$ and $y \in \Gamma_{n-r}$, by (3.5) we clearly have

$$\eta_n(\gamma \otimes q^k)(x \times y, t) = \eta_n(\gamma \otimes q^k)(x, t) \eta_n(\gamma \otimes q^k)(y, t).$$

This immediately implies the formula. □

A similar argument gives that

$$\varepsilon_n(\gamma \otimes q^k)(x, t) = (-1)^n \prod_{c \in \Gamma_*} (-\gamma(c))^{l(\rho(c))} t^{nk}, \tag{6.3}$$

where x is any element in the conjugacy class of type $\rho = (\rho(c))_{c \in \Gamma^*}$.

Formula (6.2) is equivalent to the following:

$$\eta_n(\gamma \otimes q^k)(c_\rho, t) = \prod_{c \in \Gamma_*} \prod_{i \geq 1} (\gamma \otimes q^k)(c, t^i)^{m_i(\rho(c))}. \tag{6.4}$$

The following result allows us to extend the map from $\gamma \in \Gamma^*$ to $R(\Gamma_n)$.

Proposition 6.2. *For any* $\gamma \in R(\Gamma)$*, we have*

$$\sum_{n\geq 0} ch(\eta_n(\gamma\otimes q^k))z^n = \exp\left(\sum_{n\geq 1}\frac{1}{n}\,a_{-n}(\gamma)(q^{-k}z)^n\right), \tag{6.5}$$

$$\sum_{n\geq 0} ch(\varepsilon_n(\gamma\otimes q^k))z^n = \exp\left(\sum_{n\geq 1}(-1)^{n-1}\frac{1}{n}\,a_{-n}(\gamma)(q^{-k}z)^n\right). \tag{6.6}$$

Proof. It follows from definition of ch (6.1) and (6.4) that

$$\begin{aligned}
&\sum_{n\geq 0}\mathrm{ch}(\eta_n(\gamma\otimes q^k))z^n\\
=\ &\sum_{\rho} Z_\rho^{-1}\prod_{c\in\Gamma_*}\prod_{i\geq 1} S(\gamma_{q^{ik}}(c)^{m_i(\rho(c))})a_{-\rho(c)}z^{||\rho||}q^{-||\rho||}\\
=\ &\sum_{\rho} Z_\rho^{-1}\prod_{c\in\Gamma_*}\gamma(c)^{l(\rho(c))}a_{-\rho(c)}(q^{-k}z)^{||\rho||}\\
=\ &\prod_{c\in\Gamma_*}\Big(\sum_{\lambda}(\zeta_c^{-1}\gamma(c))^{l(\lambda)}z_\lambda^{-1}a_{-\lambda}(c)(q^{-k}z)^{|\lambda|}\Big)\\
=\ &\exp\left(\sum_{n\geq 1}\frac{1}{n}\sum_{c\in\Gamma_*}\zeta_c^{-1}\gamma(c)a_{-n}(c)(q^{-k}z)^n\right)\\
=\ &\exp\left(\sum_{n\geq 1}\frac{1}{n}\,a_{-n}(\gamma)(q^{-k}z)^n\right).
\end{aligned}$$

Similarly we can prove (6.6) using the following identity

$$\begin{aligned}
\varepsilon_n(\gamma\otimes q^k)(x) &= (-1)^n\prod_{c\in\Gamma_*}\prod_{i\geq 1}(-\gamma_{q^{ik}}(c))^{m_i(\rho(c))}\\
&= (-q^k)^n\prod_{c\in\Gamma_*}\prod_{i\geq 1}(-\gamma(c))^{m_i(\rho(c))}\\
&= \varepsilon_n(\gamma)(x)q^{nk}.
\end{aligned}$$

The same argument as in the classical case (cf. [FJW]) by using (3.6) and (3.7) will show that the proposition holds for linear combination of simple characters such as $\gamma\otimes q^k - \beta\otimes q^k$, and thus it is true for any element $\gamma\otimes q^k$, where $\gamma\in R(\Gamma)$. ☐

Comparing components we obtain

$$\begin{aligned} \mathrm{ch}(\eta_n(\gamma \otimes q^k)) &= \sum_{\lambda} \frac{q^{-nk}}{z_\lambda} a_{-\lambda}(\gamma), \\ \mathrm{ch}(\varepsilon_n(\gamma \otimes q^k)) &= \sum_{\lambda} \frac{q^{-nk}}{z_\lambda} (-1)^{|\lambda|-l(\lambda)} a_{-\lambda}(\gamma), \end{aligned}$$

where the sum runs over all partitions λ of n.

Corollary 6.2. *The formula (6.4) remains valid when $\gamma \otimes q^k$ is replaced by any element $\xi \in R(\Gamma \times \mathbb{C}^\times)$. In particular $\eta_n(\xi)$ is self-dual provided that ξ is invariant under the antipode S.*

6.2. **Isometry between $R_{\Gamma\times\mathbb{C}^\times}$ and $S_{\Gamma\times\mathbb{C}^\times}$.** The symmetric algebra $S_{\Gamma\times\mathbb{C}^\times} = S_\Gamma \otimes \mathbb{C}[q, q^{-1}]$ has the following Hopf algebra structure over $\mathbb{C}$. The multiplication is the usual one, and the comultiplication is given by

$$\begin{gathered} \Delta(q^k) = q^k \otimes q^k \\ \Delta(a_n(\gamma \otimes q^k)) = a_n(\gamma \otimes q^k) \otimes q^{nk} + q^{nk} \otimes a_n(\gamma \otimes q^k), \end{gathered}$$

where $\gamma \in \Gamma^*$. The last formula is equivalent to the following:

$$\Delta(a_n(c \otimes q^k)) = a_n(c \otimes q^k) \otimes q^{nk} + q^{nk} \otimes a_n(c \otimes q^k), \tag{6.7}$$

where $c \in \Gamma_*$. The antipode is given by

$$\begin{gathered} S(q^k) = q^{-k}, \\ S(a_n(\gamma \otimes q^k)) = -a_n(\gamma \otimes q^{-k}) \end{gathered}$$

The antipode commutes with the adjoint (dual) map $*$:

$$*^2 = S^2 = Id, \qquad S* = *S. \tag{6.8}$$

Recall that we have defined a Hopf algebra structure on $R_{\Gamma\times\mathbb{C}^\times}$ in Sect. 2.

Proposition 6.3. *The characteristic map $ch : R_{\Gamma\times\mathbb{C}^\times} \longrightarrow S_{\Gamma\times\mathbb{C}^\times}$ is an isomorphism of Hopf algebras.*

Proof. It follows immediately from the definition of the comultiplication in the both Hopf algebras (cf. (2.4) and (6.7)). □

Remark 6.3. The comultiplication (6.7) is in fact induced from that of the classical case in [FJW] and only works for $C = 1$.

Remark 6.4. There is another coproduct called Drinfeld comultiplication Δ_D on the algebra $S_{\Gamma\times\mathbb{C}^\times}$ adjoined by a central element q^c. The formula on $S_{\Gamma\times\mathbb{C}^\times}$ at level c is as follows [J2]:

$$\Delta_D(a_n(\gamma)) = a_n(\gamma)\otimes q^{|n|c/2} + q^{-|n|c/2}\otimes a_n(\gamma). \tag{6.9}$$

We do not know a conceptual interpretation of the Drinfeld comultiplication in $R_{\Gamma\times\mathbb{C}^\times}$.

Recall that we have defined a bilinear form $\langle\ ,\ \rangle_\xi$ on $R_{\Gamma\times\mathbb{C}^\times}$ and a bilinear form on $S_{\Gamma\times\mathbb{C}^\times}$ denoted by $\langle\ ,\ \rangle'_\xi$, where ξ is a self-dual class function. The following lemma is immediate from our definition of $\langle\ ,\ \rangle'_\xi$ and the comultiplication Δ.

Lemma 6.5. *The bilinear form $\langle\ ,\ \rangle'_\xi$ on $S_{\Gamma\times\mathbb{C}^\times}$ can be characterized by the following two properties:*

1). $\langle a_{-n}(\beta\otimes q^k), a_{-m}(\gamma\otimes q^l)\rangle'_\xi = \delta_{n,m}q^{n(l-k)}\langle\beta,\gamma\rangle'_\xi$, $\beta,\gamma\in\Gamma^*$, $k,l\in\mathbb{Z}$.

2). $\langle fg, h\rangle'_\xi = \langle f\otimes g, \Delta h\rangle'_\xi$, where $f,g,h\in S_{\Gamma\times\mathbb{C}^\times}$, and the bilinear form on $S_{\Gamma\times\mathbb{C}^\times}\otimes S_{\Gamma\times\mathbb{C}^\times}$, is induced from $\langle\ ,\ \rangle'_\xi$ on $S_{\Gamma\times\mathbb{C}^\times}$.

Theorem 6.6. *The characteristic map is an isometry from the space $(R_{\Gamma\times\mathbb{C}^\times}, \langle\ ,\ \rangle_\xi)$ to the space $(S_{\Gamma\times\mathbb{C}^\times}, \langle\ ,\ \rangle'_\xi)$.*

Proof. By Corollary 6.2, the character value of $\eta_n(\xi)$ at an element x of type ρ is

$$\eta_n(\xi)(x) = \prod_{c\in\Gamma_*}\prod_{i\geq 1}\xi_{q^i}(c)^{m_i(\rho(c))}.$$

Thus it follows from definition that

$$\begin{aligned}\langle\sigma_{\rho\otimes q^k}, \sigma_{\rho'\otimes q^l}\rangle_\xi &= \sum_{\mu\in\mathcal{P}_n(\Gamma_*)} Z_\mu^{-1}q^{n(l-k)}\xi_q(c_\mu)\sigma_\rho(c_\mu)\sigma_{\rho'}(c_\mu)\\ &= \delta_{\rho,\rho'}Z_\rho^{-1}q^{n(l-k)}\xi(c_\rho)Z_\rho Z_\rho\\ &= \delta_{\rho,\rho'}Z_\rho q^{n(l-k)}\prod_{c\in\Gamma_*}\prod_{i\geq 1}\xi_{q^i}(c)^{m_i(\rho(c))}.\end{aligned}$$

By Lemma 6.1 and the formula (5.6), we see that

$$\langle\sigma_{\rho\otimes q^k}, \sigma_{\rho'\otimes q^l}\rangle_\xi = \langle a_{-\rho\otimes q^k}, a_{-\rho'\otimes q^l}\rangle'_\xi = \langle \mathrm{ch}(\sigma_{\rho\otimes q^k}), \mathrm{ch}(\sigma_{\rho'\otimes q^l})\rangle'_\xi.$$

Since $\sigma_{\rho\otimes q^k}, \rho\in\mathcal{P}(\Gamma_*)$ form a $\mathbb{C}$-basis of $R_{\Gamma\times\mathbb{C}^\times}$, we have shown that $\mathrm{ch}: R_{\Gamma\times\mathbb{C}^\times}\longrightarrow S_{\Gamma\times\mathbb{C}^\times}$ is an isometry. ☐

¿From now on we will not distinguish the bilinear form $\langle\ ,\ \rangle_\xi$ on $R_{\Gamma\times\mathbb{C}^\times}$ from the bilinear form $\langle\ ,\ \rangle'_\xi$ on $S_{\Gamma\times\mathbb{C}^\times}$.

7. QUANTUM VERTEX OPERATORS AND $R_{\Gamma\times\mathbb{C}^\times}$

7.1. **Vertex Operators and Heisenberg algebras in $\mathcal{F}_{\Gamma\times\mathbb{C}^\times}$.** Let Q be an integral lattice with basis α_i, $i = 0, 1, \dots, r$ endowed with a symmetric bilinear form. As in the case of $q = 1$ (cf. [FK]), we fix a 2-cocycle $\epsilon : Q \times Q \longrightarrow \mathbb{C}^\times$ such that

$$\epsilon(\alpha, \beta) = \epsilon(\beta, \alpha)(-1)^{\langle\alpha,\beta\rangle+\langle\alpha,\alpha\rangle\langle\beta,\beta\rangle}.$$

We remark that the cocycle can be constructed directly by prescribing the values of $(\alpha_i, \alpha_j) \in \{\pm 1\}$ $(i < j)$.

Let ξ be a self-dual virtual character in $R_{\Gamma\times\mathbb{C}^\times}$. Recall that the lattice $R_{\mathbb{Z}}(\Gamma)$ is a $\mathbb{Z}$-lattice under the bilinear form $\langle\ ,\ \rangle^1_\xi$, here the superscript means $q = 1$. For our purpose we will always associate a 2-cocycle ϵ as in the previous subsection to the integral lattice $(R_{\mathbb{Z}}(\Gamma), \langle\ ,\ \rangle^1_\xi)$ (and its sublattices).

Let $\mathbb{C}[R_{\mathbb{Z}}(\Gamma)]$ be the group algebra generated by e^γ, $\gamma \in R_{\mathbb{Z}}(\Gamma)$. We introduce two special operators acting on $\mathbb{C}[R_{\mathbb{Z}}(\Gamma)]$: A ($\epsilon$-twisted) multiplication operator e^α defined by

$$e^\alpha . e^\beta = \epsilon(\alpha, \beta)e^{\alpha+\beta}, \quad \alpha, \beta \in R_{\mathbb{Z}}(\Gamma),$$

and a differentiation operator ∂_α given by

$$\partial_\alpha e^\beta = \langle\alpha, \beta\rangle^1_\xi e^\beta, \quad \alpha, \beta \in R_{\mathbb{Z}}(\Gamma).$$

These two operators are then extended linearly to the space

$$\mathcal{F}_{\Gamma\times\mathbb{C}^\times} = R_{\Gamma\times\mathbb{C}^\times} \otimes \mathbb{C}[R_{\mathbb{Z}}(\Gamma)] \tag{7.1}$$

by letting them act on the $R_{\Gamma\times\mathbb{C}^\times}$ part trivially.

We define the Hopf algebra structure on $\mathbb{C}[R_{\mathbb{Z}}(\Gamma)]$ and extend the Hopf algebra structure from $R_{\Gamma\times\mathbb{C}^\times}$ to $\mathcal{F}_{\Gamma\times\mathbb{C}^\times}$ as follows.

$$\Delta(e^\alpha) = e^\alpha \otimes e^\alpha, \qquad S(e^\alpha) = e^{-\alpha}.$$

The bilinear form $\langle\ ,\ \rangle^q_\xi$ on $R_{\Gamma\times\mathbb{C}^\times}$ is extended to $\mathcal{F}_{\Gamma\times\mathbb{C}^\times}$ by

$$\langle e^\alpha, e^\beta\rangle_\xi = \delta_{\alpha,\beta}.$$

With respect to this extended bilinear form we have the $*$-action (adjoint action) on the operators e^α and ∂_α:

$$(e^\alpha)^* = e^{-\alpha}, \qquad (z^{\partial_\alpha})^* = z^{-\partial_\alpha}. \tag{7.2}$$

For each $k \in \mathbb{Z}$, we introduce the group theoretic operators $H_{\pm n}(\gamma \otimes q^k), E_{\pm n}(\gamma \otimes q^k), \gamma \in R(\Gamma), n > 0$ as the following compositions of maps:

$$\begin{aligned}
H_{-n}(\gamma \otimes q^k) \quad &: \quad R(\Gamma_m \times \mathbb{C}^\times) \xrightarrow{\eta_n(\gamma\otimes q^k)\otimes} R(\Gamma_n \times \mathbb{C}^\times) \otimes R(\Gamma_m \times \mathbb{C}^\times)\\
&\quad\ \xrightarrow{Ind\otimes m_{\mathbb{C}^\times}} R(\Gamma_{n+m} \times \mathbb{C}^\times)\\
E_{-n}(\gamma \otimes q^k) \quad &: \quad R(\Gamma_m \times \mathbb{C}^\times) \xrightarrow{\varepsilon_n(\gamma\otimes q^k)\otimes} R(\Gamma_n \times \mathbb{C}^\times) \otimes R(\Gamma_m \times \mathbb{C}^\times)\\
&\quad\ \xrightarrow{Ind\otimes m_{\mathbb{C}^\times}} R(\Gamma_{n+m} \times \mathbb{C}^\times)\\
E_{n}(\gamma \otimes q^k) \quad &: \quad R(\Gamma_m \times \mathbb{C}^\times) \xrightarrow{Res} R(\Gamma_n) \otimes R(\Gamma_{m-n} \times \mathbb{C}^\times)\\
&\quad\ \xrightarrow{\langle\varepsilon_n(\gamma\otimes q^k),\cdot\rangle_\xi} R(\Gamma_{m-n} \times \mathbb{C}^\times)\\
H_{n}(\gamma \otimes q^k) \quad &: \quad R(\Gamma_m \times \mathbb{C}^\times) \xrightarrow{Res} R(\Gamma_n) \otimes R(\Gamma_{m-n} \times \mathbb{C}^\times)\\
&\quad\ \xrightarrow{\langle\eta_n(\gamma\otimes q^k),\cdot\rangle_\xi} R(\Gamma_{m-n} \times \mathbb{C}^\times),
\end{aligned}$$

where Res and Ind are the restriction and induction functors in $R_\Gamma = \bigoplus_{n\geq 0} R(\Gamma_n)$.

We introduce their generating functions in a formal variable z:

$$\begin{aligned}
H_\pm(\gamma \otimes q^k, z) &= \sum_{n\geq 0} H_{\mp n}(\gamma \otimes q^k) z^{\pm n},\\
E_\pm(\gamma \otimes q^k, z) &= \sum_{n\geq 0} E_{\mp n}(\gamma \otimes q^k)(-z)^{\pm n}.
\end{aligned}$$

We now define the vertex operators $Y_n^\pm(\gamma \otimes q^l, k)$, $\gamma \in \Gamma^*$, $k, l \in \mathbb{Z}$, $n \in \mathbb{Z} + \langle\gamma,\gamma\rangle_\xi^1/2$ as follows.

$$\begin{aligned}
Y^+(\gamma \otimes q^l, k, z) &= \sum_{n\in\mathbb{Z}+\langle\gamma,\gamma\rangle_\xi^1/2} Y_n^+(\gamma \otimes q^l, k) z^{-n-\langle\gamma,\gamma\rangle_\xi^1/2}\\
&= H_+(\gamma \otimes q^l, z) E_-(\gamma \otimes q^{l-k}, z) e^\gamma (q^{-l}z)^{\partial_\gamma}, \qquad (7.3)
\end{aligned}$$

$$\begin{aligned}
Y^-(\gamma \otimes q^l, k, z) &= (Y^+(\gamma \otimes q^l, k, z^{-1}))^*\\
&= \sum_{n\in\mathbb{Z}+\langle\gamma,\gamma\rangle_\xi^1/2} Y_n^-(\gamma \otimes q^l, k) z^{-n-\langle\gamma,\gamma\rangle_\xi^1/2}\\
&= E_+(\gamma \otimes q^{l-k}, z) H_-(\gamma \otimes q^l, z) e^{-\gamma} (q^{-l}z)^{-\partial_\gamma}. \qquad (7.4)
\end{aligned}$$

One easily sees that the operators $Y_n^\pm(\gamma \otimes q^l, k)$ are well-defined operators acting on the space $\mathcal{F}_{\Gamma\times\mathbb{C}^\times}$.

We extend the $\mathbb{Z}_+$-gradation on $R_{\Gamma\times\mathbb{C}^\times}$ to a $\frac{1}{2}\langle\gamma,\gamma\rangle_\xi^1 + \mathbb{Z}_+$-gradation on $\mathcal{F}_{\Gamma\times\mathbb{C}^\times}$ by letting

$$\deg a_{-n}(\gamma\otimes q^k) = n, \quad \deg e^\gamma = \frac{1}{2}\langle\gamma,\gamma\rangle_\xi^1.$$

We denote by $\overline{R}_{\Gamma\times\mathbb{C}^\times}$ the subalgebra of $R_{\Gamma\times\mathbb{C}^\times}$ excluding the generators $a_n(\gamma_0)$, $n\in\mathbb{Z}^\times$. The bilinear form $\langle\ ,\ \rangle_\xi$ on

$$\overline{\mathcal{F}}_{\Gamma\times\mathbb{C}^\times} = \overline{R}_{\Gamma\times\mathbb{C}^\times}\otimes\overline{R}_{\mathbb{Z}}(\Gamma)$$

will be the restriction of $\langle\ ,\ \rangle_\xi$ on $\mathcal{F}_{\Gamma\times\mathbb{C}^\times}$ to $\overline{\mathcal{F}}_{\Gamma\times\mathbb{C}^\times}$. In the case of the second choice of ξ and $p=q^{\pm1}$, the Fock space $\overline{\mathcal{F}}_{\Gamma\times\mathbb{C}^\times}$ can also be obtained as the quotient of $\mathcal{F}_{\Gamma\times\mathbb{C}^\times}$ modulo the radical of $\langle\ ,\ \rangle_\xi$.

We define $\widetilde{a}_{-n}(\gamma\otimes q^k), n>0$ to be a map from $R_{\Gamma\times\mathbb{C}^\times}$ to itself by the following composition

$$\begin{aligned} R(\Gamma_m\times\mathbb{C}^\times) &\xrightarrow{\sigma_n(\gamma\otimes q^k)\otimes} R(\Gamma_n\times\mathbb{C}^\times)\otimes R(\Gamma_m\times\mathbb{C}^\times) \\ &\xrightarrow{Ind\otimes m_{\mathbb{C}^\times}} R(\Gamma_{n+m}\times\mathbb{C}^\times). \end{aligned}$$

We also define $\widetilde{a}_n(\gamma\otimes q^k), n>0$ to be a map from $R_{\Gamma\times\mathbb{C}^\times}$ to itself as the composition

$$\begin{aligned} R(\Gamma_m\times\mathbb{C}^\times) &\xrightarrow{Res\otimes 1} R(\Gamma_n\times\mathbb{C}^\times)\otimes R(\Gamma_{m-n}\times\mathbb{C}^\times) \\ &\xrightarrow{\langle\sigma_n(\gamma\otimes q^k),\cdot\rangle_\xi^q} R(\Gamma_{m-n}\times\mathbb{C}^\times). \end{aligned}$$

Proposition 7.1. *The operators $\widetilde{a}_n(\gamma)$, $\gamma\in\Gamma^*, n\in\mathbb{Z}^\times$ satisfy the Heisenberg algebra relations (5.1) with $C=1$.*

Proof. This is similarly proved as for the classical setting in [W]. ☐

7.2. **Group theoretic interpretation of vertex operators.** To compare the vertex operators $Y^\pm(\gamma\otimes q^l, k, z)$ with the familiar vertex operators acting in the Fock space we introduce the space

$$V_{\Gamma\times\mathbb{C}^\times} = S_{\Gamma\times\mathbb{C}^\times}\otimes\mathbb{C}[R_{\mathbb{Z}}(\Gamma)].$$

We extend the bilinear form $\langle\ ,\ \rangle_\xi^q$ in $S_{\Gamma\times\mathbb{C}^\times}$ to the space $V_{\Gamma\times\mathbb{C}^\times}$ and also extend the $\mathbb{Z}_+$-gradation on $S_{\Gamma\times\mathbb{C}^\times}$ to a $\frac{1}{2}\mathbb{Z}_+$-gradation on V_Γ.

We extend the characteristic map to the map

$$ch : \mathcal{F}_{\Gamma\times\mathbb{C}^\times}\longrightarrow V_{\Gamma\times\mathbb{C}^\times}$$

by identity on $R_{\mathbb{Z}}(\Gamma)$. Then Proposition 6.3 and Theorem 6.6 imply that we have an isometric isomorphism of Hopf algebras. We can now identify the operators from the previous subsections with the operators constructed from the Heisenberg algebra.

Theorem 7.1. *For any $\gamma \in R(\Gamma)$ and $k \in \mathbb{Z}$, we have*

$$ch\big(H_+(\gamma \otimes q^k, z)\big) = \exp\Bigg(\sum_{n\geq 1} \frac{1}{n}\, a_{-n}(\gamma)(q^{-k}z)^n\Bigg), \tag{7.5}$$

$$ch\big(E_+(\gamma \otimes q^k, z)\big) = \exp\Bigg(-\sum_{n\geq 1} \frac{1}{n}\, a_{-n}(\gamma)(q^{-k}z)^n\Bigg), \tag{7.6}$$

$$ch\big(H_-(\gamma \otimes q^k, z)\big) = \exp\Bigg(\sum_{n\geq 1} \frac{1}{n}\, a_{n}(\gamma)(q^{-k}z)^{-n}\Bigg), \tag{7.7}$$

$$ch\big(E_-(\gamma \otimes q^k, z)\big) = \exp\Bigg(-\sum_{n\geq 1} \frac{1}{n}\, a_{n}(\gamma)(q^{-k}z)^{-n}\Bigg). \tag{7.8}$$

Proof. The first and second identities were essentially established in Proposition 6.2 together with Lemma 6.1, where the components are viewed as operators acting on $R_{\Gamma\times\mathbb{C}^\times}$ or $S_{\Gamma\times\mathbb{C}^\times}$. Note that $a_n(\gamma \otimes q^k) = a_n(\gamma)q^{kn}$.

We observe from definition that the adjoint $*$-action of $E_+(\gamma\otimes q^k, z)$ and $H_-(\gamma \otimes q^k, z)$ with respect to the bilinear form $\langle\ ,\ \rangle_\xi^q$ are $E_-(\gamma \otimes q^k, z^{-1})$ and $H_-(\gamma \otimes q^k, z^{-1})$ respectively. The third and fourth identities are obtained by applying the adjoint action $*$ to the first two identities. □

Remark 7.2. Replacing γ by $-\gamma$ in (7.5) and (7.7) we obtain the equivalent formulas (7.6) and (7.8) respectively.

Applying the characteristic map to the vertex operators $Y^{\pm}(\gamma, k, z)$, we obtain the following group theoretical explanation of vertex operators acting on the Fock space $\mathcal{F}_{\Gamma\times\mathbb{C}^\times}$.

Theorem 7.3. *For any $\gamma \in R_\Gamma$ and $k \in \mathbb{Z}$, we have*

$$\begin{aligned} &Y^+(\gamma, k, z)\\ &= \exp\Bigg(\sum_{n\geq 1}\frac{1}{n}\,\widetilde{a}_{-n}(\gamma)z^n\Bigg)\ \exp\Bigg(-\sum_{n\geq 1}\frac{1}{n}\,\widetilde{a}_{n}(\gamma)q^{-kn}z^{-n}\Bigg)e^{\gamma}z^{\partial_\gamma}\\ &= ch(H_+(\gamma, z))ch(S(H_+(\gamma\otimes q^k, z^{-1})^*))e^{\gamma}z^{\partial_\gamma}, \end{aligned}$$

$$\begin{aligned} &Y^-(\gamma, k, z)\\ &= \exp\Bigg(-\sum_{n\geq 1}\frac{1}{n}\,\widetilde{a}_{-n}(\gamma)q^{kn}z^n\Bigg)\ \exp\Bigg(\sum_{n\geq 1}\frac{1}{n}\,\widetilde{a}_{n}(\gamma)z^{-n}\Bigg)e^{-\gamma}z^{-\partial_\gamma}\\ &= ch(S(H_+(\gamma\otimes q^k, z^{-1})))ch(H_+(\gamma, z)^*)e^{-\gamma}z^{-\partial_\gamma}. \end{aligned}$$

We note that for $\gamma \in \Gamma^*, l \in \mathbb{Z}$

$$Y^{\pm}(\gamma \otimes q^l, k, z) = Y^{\pm}(\gamma, k, q^{-l}z). \tag{7.9}$$

It follows from Theorem 7.3 that

$$\begin{aligned} ch\big(Y^{\pm}(\gamma, k, z)\big) &= X^{\pm}(\gamma, k, z) \\ &= \exp\Big(-\sum_{n\geq 1} \frac{1}{n}\, a_{-n}(\gamma) q^{n(k\mp k)/2} z^n\Big) \\ &\quad \times \exp\Big(\sum_{n\geq 1} \frac{1}{n}\, a_n(\gamma) q^{n(-k\mp k)/2} z^{-n}\Big) e^{\pm\gamma} z^{\pm\partial_\gamma}. \end{aligned}$$

In general the vertex operators $Y^{\pm}(\gamma, k, z)$ (for $k \in \mathbb{Z}$) generalize the vertex operators considered in [J3] (for $k = \pm 1$). When $q = 1$ they specialize to the vertex operators $Y^{\pm}(\gamma, z)$ studied in [FJW].

8. Basic representations and the McKay correspondence

8.1. **Quantum toroidal algebras.** Let Q be the root lattice of an affine Lie algebra of simply laced type A, D, or E with the invariant form $(\ |\ )$. The quantum toroidal algebra $U_q(\widehat{\widehat{\mathfrak{g}}})$ is the associative algebra generated by $x_i^{\pm}(n)$, $a_i(m)$, q^d, q^c , $0 \leq i \leq r, n, m \in \mathbb{Z}$ subject to the following relations [GKV]:

$$q^d a_i(n) q^{-d} = q^n a_i(n), q^d x_i^{\pm}(n) q^{-d} = q^n x_i^{\pm}(n), \tag{8.1}$$

$$[a_i(m), a_j(n)] = \delta_{m,-n} \frac{[(\alpha_i|\alpha_j)m]}{m} \frac{q^{mc} - q^{-mc}}{q - q^{-1}}, \tag{8.2}$$

$$[a_i(m), x_j^{\pm}(n)] = \pm \frac{[(\alpha_i|\alpha_j)m]}{m} q^{\mp|m|c/2} x_j^{\pm}(m+n), \tag{8.3}$$

$$(z - q^{\pm(\alpha_i,\alpha_j)}w) x_i^{\pm}(z) x_j^{\pm}(w) = x_j^{\pm}(w) x_i^{\pm}(z)(q^{\pm(\alpha_i,\alpha_j)}z - w), \tag{8.4}$$

(8.5)

$$[x_i^{+}(z), x_j^{-}(w)] = \frac{\delta_{ij}\{\delta(zw^{-1}q^{-c})\psi_i^{+}(wq^{c/2}) - \delta(zw^{-1}q^{c})\psi_i^{-}(zq^{c/2})\}}{q - q^{-1}},$$

$$\begin{aligned} Sym_{z_1,\ldots z_N} \sum_{s=0}^{N=1-(\alpha_i,\alpha_j)} (-1)^s \begin{bmatrix} N \\ s \end{bmatrix} x_i^{\pm}(z_1) \cdots x_i^{\pm}(z_s) \cdot \\ \cdot x_j^{\pm}(w) x_i^{\pm}(z_{s+1}) \cdots x_i^{\pm}(z_N) = 0, \quad \text{for} \quad (\alpha_i|\alpha_j) \leq 0, \end{aligned} \tag{8.6}$$

where the generators $\alpha(n)$ are related to $\psi_i^{\pm}(\pm n)$ via:

$$\psi_i^{\pm}(z) = \sum_{n\geq 0} \psi_i^{\pm}(\pm n) z^{\mp n} = k_i^{\pm 1} exp(\pm(q - q^{-1}) \sum_{n>0} \alpha_i(\pm n) z^{\mp n}), \tag{8.7}$$

and the Gaussian polynomial

$$\begin{bmatrix} m \\ n \end{bmatrix} = \frac{[m]!}{[n]![m-n]!}, \qquad [n]! = [n][n-1]\cdots[1].$$

The generating function of $x_n^{\pm}$ are defined by

$$x_i^{\pm}(z) = \sum_{n\in\mathbb{Z}} x_i^{\pm}(n) z^{-n-1}, \qquad i = 0, \dots, r.$$

The quantum toroidal algebra contains a special subalgebra– the quantum affine algebra $U_q(\widehat{\mathfrak{g}})$, which is generated by simply omitting the generators associated to $i = 0$. The relations are called the Drinfeld realization of the quantum affine algebras.

In the case of type A, the quantum toroidal algebra $U_q(\widehat{\widehat{\mathfrak{g}}})$ admits a further deformation $U_{q,p}(\widehat{\widehat{\mathfrak{g}}})$. Let (b_{ij}) be the skew-symmetric $(r+1) \times (r+1)$-matrix

$$\begin{pmatrix} 0 & 1 & 0 & \cdots & 0 & -1 \\ -1 & 0 & 1 & \cdots & 0 & 0 \\ 0 & -1 & 0 & \cdots & 0 & 0 \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & 0 & \cdots & -1 & 0 \end{pmatrix}. \tag{8.8}$$

The quantum toroidal algebra $U_{q,p}(\widehat{\widehat{\mathfrak{g}}})$ is the associative algebra generated by $x_{in}^{\pm}$, $a_i(m)$, q^{d_1}, q^{d_2}, q^c , $0 \le i \le r, m, n \in \mathbb{Z}$ subject to the following relations [GKV, VV]:

$$q^{d_1} a_i(n) q^{-d_1} = q^n a_i(n), q^{d_1} x_i^{\pm}(n) q^{-d_1} = q^n x_i^{\pm}(n), \tag{8.9}$$

$$q^{d_2} a_i(n) q^{-d_2} = a_i(n) \tag{8.10}$$

$$q^{d_2} x_i^{\pm}(n) q^{-d_2} = q^{\pm\delta_{n0}} x_i^{\pm}(n), \tag{8.11}$$

$$[a_i(m), a_j(n)] = \delta_{m,-n} \frac{[(\alpha_i|\alpha_j)m]}{m} \frac{q^{mc} - q^{-mc}}{q - q^{-1}} p^{mb_{ij}}, \tag{8.12}$$

$$[a_i(m), x_j^{\pm}(n)] = \pm \frac{[(\alpha_i|\alpha_j)m]}{m} q^{\mp|m|c/2} p^{mb_{ij}} x_j^{\pm}(m+n), \tag{8.13}$$

(8.14)
$$(p^{b_{ij}} z - q^{\pm(\alpha_i|\alpha_j)} w) x_i^{\pm}(z) x_j^{\pm}(w) = x_j^{\pm}(w) x_i^{\pm}(z) (p^{b_{ij}} q^{\pm(\alpha_i|\alpha_j)} z - w),$$

(8.15)
$$[x_i^{+}(z), x_j^{-}(w)] = \frac{\delta_{ij}\{\delta(zw^{-1}q^{-c})\psi_i^{+}(wq^{c/2}) - \delta(zw^{-1}q^{c})\psi_i^{-}(zq^{c/2})\}}{q - q^{-1}},$$

$$\begin{aligned} Sym_{z_1,\dots z_N} \sum_{s=0}^{N=1-(\alpha_i|\alpha_j)} (-1)^s \begin{bmatrix} N \\ s \end{bmatrix} x_i^{\pm}(z_1) \cdots x_i^{\pm}(z_s) \cdot \\ \cdot x_j^{\pm}(w) x_i^{\pm}(z_{s+1}) \cdots x_i^{\pm}(z_N) = 0, \quad \text{for} \quad (\alpha_i|\alpha_j) \leq 0, \end{aligned} \tag{8.16}$$

where the generators $a_i(n)$ are related to $\psi_i^{\pm}(\pm m)$ via:

(8.17)
$$\psi_i^{\pm}(z) = \sum_{n \geq 0} \psi_i^{\pm}(\pm n) z^{\mp n} = k_i^{\pm 1} exp(\pm(q - q^{-1}) \sum_{n > 0} \alpha_i(\pm n) z^{\mp n}).$$

We recall that the *basic module* of $U_q(\widehat{\mathfrak{g}})$ is the simple module generated by the highest weight vector v_0 such that

$$a_i(n+1).v_0 = 0, \qquad x_i^{\pm}(n).v_0 = 0, \qquad n \geq 0$$
$$q^c.v_0 = qv_0, \qquad q^d.v_0 = v_0.$$

We say a module is of level one if q^c acts as q.

8.2. **A new form of McKay correspondence.** In this subsection we let Γ to be a finite subgroup of SU_2 and consider two distinguished choices of the class function ξ in $R_{\Gamma \times \mathbb{C}^\times}$ introduced in Sect. 4.2.

First we consider

$$\xi = \gamma_0 \otimes (q + q^{-1}) - \pi \otimes 1_{\mathbb{C}^\times},$$

where π is the character of the two-dimensional natural representation of Γ in SU_2.

The Heisenberg algebra in this case has the following relations (cf. Prop. 5.1 and (4.2)).

$$[a_m(\gamma_i), a_n(\gamma_j)] = \begin{cases} m\delta_{m,-n}(q^m + q^{-m})C, & i = j \\ m\delta_{m,-n}a^1_{ij}C, & i \neq j \end{cases}, \tag{8.18}$$

where a^1_{ij} are the entries of the affine Cartan matrix of ADE type (see (3.4) at $d = 2$).

When $\Gamma \neq \mathbb{Z}/2\mathbb{Z}$ or 1, the relations (8.18) can be simply written as follows:

$$[a_m(\gamma_i), a_n(\gamma_j)] = m\delta_{m,-n}[a_{ij}]_{q^m}C.$$

Recall that the matrix $A^1 = (\langle\gamma_i, \gamma_j\rangle^1_\xi) = (a^1_{ij})_{0\leq i,j\leq r}$ is the Cartan matrix for the corresponding affine Lie algebra [Mc]. In particular $a^1_{ii} = 2$; $a^1_{ij} = 0$ or -1 when $i \neq j$ and $\Gamma \neq \mathbb{Z}/2\mathbb{Z}$. In the case of $\Gamma = \mathbb{Z}/2\mathbb{Z}$, $a^1_{01} = a^1_{10} = -2$. Let $\mathfrak{g}$ (resp. $\hat{\mathfrak{g}}$) be the corresponding simple Lie algebra (resp. affine Lie algebra) associated to the Cartan matrix $(a^1_{ij})_{1\leq i,j\leq r}$ (resp. A). Note that the lattice $R_{\mathbb{Z}}(\Gamma)$ is even in this case.

We define the normal ordered product of vertex operators as follows.

$$\begin{aligned}
&: Y^+(\gamma_i, k, z)Y^+(\gamma_j, k', w) : \\
=&H_+(\gamma_i, z)H(\gamma_j, w)S(H_+(\gamma_i \otimes q^k, z^{-1})^* H_+(\gamma_j \otimes q^{k'}, w^{-1})^*) \\
&\times e^{\gamma_i+\gamma_j} z^{\partial_{\gamma_i}} w^{\partial_{\gamma_j}}, \\
&: Y^+(\gamma_i, k, z)Y^-(\gamma_j, k', w) : \\
=&H_+(\gamma_i, z)H(-\gamma_j \otimes q^{-k'}, w)S(H_+(\gamma_i \otimes q^k, z^{-1})^* H_+(-\gamma_j \otimes q^{k'}, w^{-1})^*) \\
&\times e^{\gamma_i-\gamma_j} z^{\partial_{\gamma_i}} w^{-\partial_{\gamma_j}}.
\end{aligned}$$

Other normal ordered products are defined similarly.

We introduce for $a \in \mathbb{R}$ the following q-function:

$$\begin{aligned}
(1 - z)^a_{q^2} &= \frac{(q^{-a+1}z; q^2)_\infty}{(q^{a+1}z; q^2)_\infty} = exp\Big(-\sum_{n=1}^{\infty} \frac{[an]}{n[n]} z^n\Big) \\
&= \sum_{m=0}^{\infty} \begin{bmatrix} a \\ m \end{bmatrix} (-z)^m,
\end{aligned} \tag{8.19}$$

where we expand the power series using the q-binomial theorem and

$$\begin{bmatrix} a \\ m \end{bmatrix} = \frac{(q^a - q^{-a})(q^{a-1} - q^{-a+1}) \cdots (q^{a-m+1} - q^{-a+m-1})}{(q^m - q^{-m})(q^{m-1} - q^{-m+1}) \cdots (q - q^{-1})},$$

$$(a; q)_\infty = \prod_{n=0}^{\infty} (1 - aq^n).$$

When a is a non-negative integer, $\begin{bmatrix} a \\ m \end{bmatrix}$ equals the Gaussian polynomial.

The identities in the following theorems are understood as usual by means of correlation functions (cf. e.g. [FJ, J1]).

Theorem 8.1. *Let* $\xi = \gamma_0 \otimes (q + q^{-1}) - \pi \otimes 1_{\mathbb{C}^\times}$. *Then the vertex operators* $Y^\pm(\gamma_i, k, z), Y^\pm(-\gamma_j, k, z)$, $\gamma_i \in \Gamma^*, k \in \mathbb{Z}$ *acting on the group theoretically defined Fock space* $\mathcal{F}_{\Gamma \times \mathbb{C}^\times}$ *satisfy the following relations.*

$$Y^\pm(\gamma_i, k, z) Y^\pm(\gamma_j, k, w) = \epsilon(\gamma_i, \gamma_j) : Y^\pm(\gamma_i, k, z) Y^\pm(\gamma_j, k, w) :$$
$$\times \begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ (z - q^{\mp k} w)^{-1} & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - q^{\mp k - 1} w)(z - q^{\mp k + 1} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases},$$

$$Y^\pm(\gamma_i, k, z) Y^\mp(\gamma_j, k, w) = \epsilon(\gamma_i, \gamma_j) : Y^\pm(\gamma_i, k, z) Y^\mp(\gamma_j, k, w) :$$
$$\times \begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ (z - w)^{-1} & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - qw)(z - q^{-1} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases},$$

$$\begin{aligned} & Y^\pm(\gamma_i, k, z) Y^\pm(-\gamma_j, -k, w) \\ & \quad = \epsilon(\gamma_i, \gamma_j) : Y^\pm(\gamma_i, k, z) Y^\pm(-\gamma_j, -k, w) : \\ & \quad \times \begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ (z - q^{\mp k} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - q^{\mp k - 1} w)^{-1} (z - q^{\mp k + 1} w)^{-1} & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases}, \end{aligned}$$

$$\begin{aligned} & Y^+(\gamma_i, k, z) Y^-(-\gamma_j, -k, w) \\ & \quad = \epsilon(\gamma_i, \gamma_j) : Y^+(\gamma_i, k, z) Y^-(-\gamma_j, -k, w) : \\ & \quad \times \begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ (z - q^{-2k} w)^{-1} & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - q^{-2k - 1} w)(z - q^{-2k + 1} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases}, \end{aligned}$$

$$\begin{aligned} &Y^-(\gamma_i, k, z)Y^+(-\gamma_j, -k, w) \\ &\qquad = \epsilon(\gamma_i, \gamma_j) : Y^-(\gamma_i, k, z)Y^+(-\gamma_j, -k, w) : \\ &\qquad \begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ (z-w)^{-1} & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z-qw)(z-q^{-1}w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases} . \end{aligned}$$

Proof. It is a routine computation to see that:

$$\begin{aligned} & E_-(\gamma_i \otimes q^k, z)H_+(\gamma_j \otimes q^l, w) \\ = \quad & H_+(\gamma_j \otimes q^l, w)E_-(\gamma_i \otimes q^k, z)(1 - \frac{w}{z}q^{l-k})_{q^2}^{\langle \gamma_i, \gamma_j \rangle_\xi^1}, \end{aligned}$$

where the q-analog of the power series $(1-x)_{q^2}^n$ is defined in (8.19).

In particular, we have

$$\begin{aligned} (1 - w/z)_{q^2} &= 1 - w/z, \\ (1 - w/z)_{q^2}^2 &= (1 - qw/z)(1 - q^{-1}w/z). \end{aligned}$$

Then the theorem is proved by observing that $z^\gamma e^{\partial_\beta} = z^{\langle \gamma, \beta \rangle_\xi^1} e^{\partial_\beta} z^\gamma$. ☐

Remark 8.2. Replacing the vertex operator $Y^\pm$ by $X^\pm$ via the characteristic map *ch* in the above formulas, we get the corresponding formulas for vertex operators $X^\pm(\gamma, k, z)$ acting on $V_{\Gamma \times \mathbb{C}^\times}$.

Now we consider the second distinguished class function

$$\xi^{q,p} = \gamma_0 \otimes (q + q^{-1}) - (\gamma_1 \otimes p + \gamma_r \otimes p^{-1}),$$

when Γ is a cyclic group of order $r+1$.

In this case the Heisenberg algebra (5.3) has the following relations according to Prop. 5.1 and (4.3):

$$[a_m(\gamma_i), a_n(\gamma_j)] = m\delta_{m,-n}[a_{ij}^1]_{q^m} p^{mb_{ij}} C, \tag{8.20}$$

where a_{ij}^1 are the entries of the affine Cartan matrix of type A and $r \geq 2$. This is the same Heisenberg subalgebra ($c = 1$) in $U_{q,p}(\widehat{\mathfrak{g}})$ provided that we identify

$$a_i(n) = \frac{[n]}{n} a_n(\gamma_i).$$

Recall that (b_{ij}) is the skew-symmetric matrix given in (8.8). We need to slightly modify the definition of the middle term in the vertex operators. For each $i = 0, 1, \ldots, r$ we define the modified operator $z^{\partial_{\gamma,p}}$ on the group algebra $\mathbb{C}[R_\mathbb{Z}(\Gamma)]$ by

$$z^{\partial_{\gamma_i,p}} e^\beta = z^{\langle \gamma_i, \beta \rangle_\xi^1} p^{-\frac{1}{2}\sum_{j=1}^r \langle \gamma_i, m_j \gamma_j \rangle_\xi^1 b_{ij}} e^\beta, \tag{8.21}$$

where $\beta = \sum_j m_j \gamma_j \in R_{\mathbb{Z}}(\Gamma)$.

We then replace the operator $z^{\pm \partial_{\gamma_i}}$ in the definition of the vertex operators $Y^{\pm}(\gamma_i, k, z)$ by the operator $z^{\pm \partial_{\gamma_i, p}}$. The formulas in Theorems 7.3 remain true after the term $z^{\pm \partial}$ appearing in the formulas are modified accordingly.

The proof of the following theorem is similar to that of Theorem 8.1.

Theorem 8.3. *Let Γ be a cyclic group of order $r+1$ and let $\xi = \gamma_0 \otimes (q + q^{-1}) - (\gamma_1 \otimes p + \gamma_r \otimes p^{-1})$. The vertex operators $Y^{\pm}(\gamma_i, k, z)$ and $Y^{\pm}(-\gamma_i, k, z)$, $\gamma_i \in \Gamma^*$ acting on the group theoretically defined Fock space $\mathcal{F}_{\Gamma \times \mathbb{C}^\times}$ satisfy the following relations.*

$$Y^{\pm}(\gamma_i, k, z) Y^{\pm}(\gamma_j, k, w) = \epsilon(\gamma_i, \gamma_j) : Y^{\pm}(\gamma_i, k, z) Y^{\pm}(\gamma_j, k, w) :$$
$$\begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ p^{-\frac{1}{2} b_{ij}} (z - q^{\mp k} p^{b_{ij}} w)^{-1} & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - q^{\mp k - 1} w)(z - q^{\mp k + 1} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases},$$

$$Y^{\pm}(\gamma_i, k, z) Y^{\mp}(\gamma_j, k, w) = \epsilon(\gamma_i, \gamma_j) : Y^{\pm}(\gamma_i, k, z) Y^{\mp}(\gamma_j, k, w) :$$
$$\begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ p^{-\frac{1}{2} b_{ij}} (z - p^{b_{ij}} w)^{-1} & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - q w)(z - q^{-1} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases},$$

$$\begin{aligned} & Y^{\pm}(\gamma_i, k, z) Y^{\pm}(-\gamma_j, -k, w) \\ & \quad = \epsilon(\gamma_i, \gamma_j) : Y^{\pm}(\gamma_i, k, z) Y^{\pm}(-\gamma_j, -k, w) : \\ & \quad \begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ p^{-\frac{1}{2} b_{ij}} (z - q^{\mp k} p^{b_{ij}} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - q^{\mp k - 1} w)^{-1} (z - q^{\mp k + 1} w)^{-1} & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases}, \end{aligned}$$

$$\begin{aligned} & Y^{+}(\gamma_i, k, z) Y^{-}(-\gamma_j, -k, w) \\ & \quad = \epsilon(\gamma_i, \gamma_j) : Y^{+}(\gamma_i, k, z) Y^{-}(-\gamma_j, -k, w) : \\ & \quad \begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ p^{-\frac{1}{2} b_{ij}} (z - q^{-2k} p^{b_{ij}} w)^{-1} & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - q^{-2k - 1} w)(z - q^{-2k + 1} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases}, \end{aligned}$$

$$\begin{aligned} & Y^{-}(\gamma_i, k, z) Y^{+}(-\gamma_j, -k, w) \\ & \quad = \epsilon(\gamma_i, \gamma_j) : Y^{-}(\gamma_i, k, z) Y^{+}(-\gamma_j, -k, w) : \\ & \quad \begin{cases} 1 & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 0 \\ p^{-\frac{1}{2} b_{ij}} (z - p^{b_{ij}} w)^{-1} & \langle \gamma_i, \gamma_j \rangle_\xi^1 = -1 \\ (z - q w)(z - q^{-1} w) & \langle \gamma_i, \gamma_j \rangle_\xi^1 = 2 \end{cases}. \end{aligned}$$

Remark 8.4. Replacing the vertex operators $Y^{\pm}$ by $X^{\pm}$ via the characteristic map *ch* we obtain the corresponding results on the space $V_{\Gamma\times\mathbb{C}^\times}$.

8.3. **Quantum vertex representations of $U_q(\widehat{\mathfrak{g}})$.** For each $i = 0, \dots, r$ let

$$\widetilde{a}_i(n) = \frac{[n]}{n} a_n(\gamma_i).$$

It follows from (5.3) and (8.18) that

$$[\widetilde{a}_i(m), \widetilde{a}_j(n)] = \delta_{m,-n} \frac{[m\langle\gamma_i, \gamma_j\rangle_\xi^1]}{m}[m]. \tag{8.22}$$

According to McKay, the bilinear form $\langle\gamma_i, \gamma_j\rangle_\xi^1$ is exactly the same as the invariant form $(\ |\ )$ of the root lattice of the affine Lie algebra $\widehat{\mathfrak{g}}$. This implies that the commutation relations (8.22) are exactly the commutation relations (8.2) of the Heisenberg algebra in $U_q(\widehat{\mathfrak{g}})$ if we identify $\widetilde{a}_i(n)$ with $a_i(n)$. Thus the Fock space $S_{\Gamma\times\mathbb{C}^\times}$ is a level one representation for the Heisenberg subalgebra in $U_q(\widehat{\mathfrak{g}})$. Under the new variable (by identifying $a_i(n)$ with $\widetilde{a}_i(n)$) and after a q-shift we obtain that

$$\begin{aligned}
&X^+(\gamma_i \otimes q^{-k/2}, k, z) \\
&= \exp\Big(\sum_{n\geq 1} \frac{a_i(-n)}{[n]} q^{kn/2} z^n\Big) \exp\Big(-\sum_{n\geq 1} \frac{a_i(n)}{[n]} q^{kn/2} z^{-n}\Big) e^{\gamma} z^{\partial_\gamma}, \\
&X^-(\gamma_i \otimes q^{-k/2}, k, z) \\
&= \exp\Big(-\sum_{n\geq 1} \frac{a_i(-n)}{[n]} q^{-kn/2} z^n\Big) \exp\Big(\sum_{n\geq 1} \frac{a_i(n)}{[n]} q^{-kn/2} z^{-n}\Big) e^{-\gamma} z^{-\partial_\gamma}.
\end{aligned}$$

The following theorem gives a q-deformation of the new form of McKay correspondence in [FJW] and provides a direct connection from a finite subgroup Γ of SU_2 to the quantum toroidal algebra $U_q(\widehat{\mathfrak{g}})$ of *ADE* type.

Theorem 8.5. *Given a finite subgroup Γ of SU_2, each of the following correspondence gives a vertex representation of the quantum toroidal algebra $U_q(\widehat{\mathfrak{g}})$ on the Fock space $\mathcal{F}_{\Gamma\times\mathbb{C}^\times}$:*

$$x_i^{\pm}(n) \longrightarrow Y_n^{\pm}(\gamma_i, -1),$$

$$a_i(n) \longrightarrow \frac{[n]}{n} a_n(\gamma_i), \qquad q^c \longrightarrow q;$$

or

$$\begin{aligned} x_i^{\pm}(n) &\longrightarrow Y_n^{\mp}(-\gamma_i, 1), \\ a_i(n) &\longrightarrow \frac{[n]}{n} a_n(\gamma_i), \qquad q^c \longrightarrow q, \end{aligned}$$

where $i = 0, \dots, r$, *and* $n \in \mathbb{Z}$.

Proof. Using the usual method of q-vertex operator calculus [FJ, J1] and Theorem 8.1 we see that the vertex operators $Y^{\pm}(\gamma_i, \pm 1, z)$ satisfy relations (8.3), (8.4) and (8.6). Observe further that the above vertex operators at $k = \pm 1$ have the same form as those in the basic representations of the quantum affine algebras (see [FJ]). Thus the relations (8.5) and (8.7) are also verified. For each fixed $k = 1$ or -1 we have shown that the operators $Y^{\pm}(\gamma_i, \pm 1, z)$ give a level one representation of the quantum toroidal algebra $U_q(\widehat{\widehat{\mathfrak{g}}})$ (see also [Sa, J3]). $\square$

Remark 8.6. Replacing $Y^{\pm}$ by $X^{\pm}$ in the above theorem, we obtain a vertex representation of $U_q(\widehat{\widehat{\mathfrak{g}}})$ in the space $V_{\Gamma \times \mathbb{C}^\times}$.

We can easily get the basic representation of the quantum affine algebra $U_q(\widehat{\mathfrak{g}})$ on a certain distinguished subspace of $\mathcal{F}_{\Gamma \times \mathbb{C}^\times}$.

Denote by $\overline{S}_{\Gamma \times \mathbb{C}^\times}$ the symmetric algebra generated by $a_{-n}(\gamma_i)$, $n > 0$, $i = 1, \dots, r$ over $\mathbb{C}[q, q^{-1}]$. $\overline{S}_{\Gamma \times \mathbb{C}^\times}$ is isometric to $\overline{R}_{\Gamma \times \mathbb{C}^\times}$.

We define

$$\overline{\mathcal{F}}_{\Gamma \times \mathbb{C}^\times} = \overline{R}_{\Gamma \times \mathbb{C}^\times} \otimes \mathbb{C}[\overline{R}_{\mathbb{Z}}(\Gamma)] \cong \overline{S}_{\Gamma \times \mathbb{C}^\times} \otimes \mathbb{C}[\overline{R}_{\mathbb{Z}}(\Gamma)].$$

The space $V_{\Gamma \times \mathbb{C}^\times}$ associated to the lattice $R_{\mathbb{Z}}(\Gamma)$ is isomorphic to the tensor product of the space $\overline{R}_{\Gamma \times \mathbb{C}^\times}$ and $R_{\mathbb{Z}}(\Gamma)$ as well as the space associated to the rank 1 lattice $\mathbb{Z}\alpha_0$.

Corollary 8.7. *Given a finite subgroup* Γ *of* SU_2, *each of the following correspondence gives the basic representation of the quantum affine algebra* $U_q(\widehat{\mathfrak{g}})$ *on the Fock space* $\overline{\mathcal{F}}_{\Gamma \times \mathbb{C}^\times}$*:*

$$\begin{aligned} x_i^{\pm}(n) &\longrightarrow Y_n^{\pm}(\gamma_i, -1), \\ a_i(n) &\longrightarrow \frac{[n]}{n} a_n(\gamma_i), \qquad q^c \longrightarrow q; \end{aligned}$$

or

$$\begin{aligned} x_i^{\pm}(n) &\longrightarrow Y_n^{\mp}(-\gamma_i, 1), \\ a_i(n) &\longrightarrow \frac{[n]}{n} a_n(\gamma_i), \qquad q^c \longrightarrow q, \end{aligned}$$

where $i = 1, \dots, r$.

In the case of our second distinguished class function

$$\xi^{q,p} = \gamma_0 \otimes (q + q^{-1}) - (\gamma_1 \otimes p + \gamma_r \otimes p^{-1}),$$

we need to consider the Fock space

$$\widetilde{\mathcal{F}}_{\Gamma\times\mathbb{C}^\times} = R_{\Gamma\times\mathbb{C}^\times} \otimes \mathbb{C}[R_{\mathbb{Z}}(\Gamma)/R^0_{\mathbb{Z}}] \cong S_{\Gamma\times\mathbb{C}^\times} \otimes \mathbb{C}[\overline{R}_{\mathbb{Z}}(\Gamma)],$$

where $R^0_{\mathbb{Z}}$ is the radical of the bilinear form $\langle\ ,\ \rangle^1_\xi$. The correspondence space for $\widetilde{\mathcal{F}}_{\Gamma\times\mathbb{C}^\times}$ under the characteristic map ch will be denoted $\widetilde{V}_{\Gamma\times\mathbb{C}^\times}$.

Using similar method as in the proof of Theorem 8.5 we derive the the following theorem.

Theorem 8.8. *Let Γ be a cyclic group of order $r+1 \geq 2$ and $p = q^{\pm 1}$. Each of the following correspondence gives the basic representation of $U_q(\widehat{\mathfrak{g}})$ on $\widetilde{\mathcal{F}}_{\Gamma\times\mathbb{C}^\times}$:*

$$\begin{aligned} x_i^{\pm}(n) &\longrightarrow Y_n^{\pm}(\gamma_i, -1), \\ a_i(n) &\longrightarrow \frac{[n]}{n} a_n(\gamma_i), \qquad q^c \longrightarrow q; \end{aligned}$$

or

$$\begin{aligned} x_i^{\pm}(n) &\longrightarrow Y_n^{\mp}(-\gamma_i, 1), \\ a_i(n) &\longrightarrow \frac{[n]}{n} a_n(\gamma_i), \qquad q^c \longrightarrow q, \end{aligned}$$

where $i = 0, \dots, r$.

Remark 8.9. The algebraic picture obtained by replacing the vertex operator $Y^{\pm}$ by $X^{\pm}$ and $\mathcal{F}_{\Gamma\times\mathbb{C}^\times}$ by $\widetilde{V}_{\Gamma\times\mathbb{C}^\times}$ in the above Theorem was given by Sato [Sa].

This theorem partly shows why the two-parameter deformation for $U_q(\widehat{\mathfrak{g}})$ is only available in the case of type A. It also singles out the special case of $p = q^{\pm 1}$, where the matrix of the bilinear form $\langle\ ,\ \rangle_\xi^{q,q^{\pm 1}}$ is semi-definite positive (see Sect. 4.2) which permits the factorization of $\mathcal{F}_{\Gamma\times\mathbb{C}^\times}$ into $\overline{\mathcal{F}}_{\Gamma\times\mathbb{C}^\times}$.

We remark that our method can be generalized by replacing $R(\Gamma)$ by any finite dimensional Hopf algebra with a Haar measure. A more general deformation is obtained by replacing $\mathbb{C}^\times$ by any torsion-free abelian group. In another direction one can replace $\mathbb{C}^\times$ by its finite analog $\mathbb{Z}/r\mathbb{Z}$ to study $U_q(\widehat{\mathfrak{g}})$ at rth roots of unity.

References

[A] E. Abe, *Hopf algebras*, Cambridge Tracts in Mathematics, **74**. Cambridge University Press, Cambridge-New York, 1980.

[CG] N. Chriss and V. Ginzburg, *Representation theory and complex geometry*, Birkhäuser, Boston, 1997.

[FJ] I. B. Frenkel and N. Jing, *Vertex representations of quantum affine algebras*, Proc. Natl. Acad. Sci. USA **85** (1988), 9373-9377.

[FJW] I. B. Frenkel, N. Jing and W. Wang, *Vertex representations via finite groups and the McKay correspondence*, Int'l. Math. Res. Notices, to appear. (math.QA/9907166)

[FK] I. B. Frenkel and V. G. Kac, *Basic representations of affine Lie algebras and dual resonance models*, Invent. Math. **62** (1980), 23–66.

[GKV] V. Ginzburg, M. Kapranov and E. Vasserot, *Langlands reciprocity for algebraic surfaces*, Math. Res. Lett. **2** (1995), 147-160.

[Gr] I. Grojnowski, *Instantons and affine algebras I: the Hilbert scheme and vertex operators*, Math. Res. Lett. **3** (1996) 275–291.

[J1] N. Jing, *Twisted vertex representations of quantum affine algebras*, Invent. Math. **102** (1990), 663-660.

[J2] N. Jing, *Higher level representations of the quantum affine algebra* $U_q(\widehat{sl}_2)$, J. Alg. **182** (1996), 448-468.

[J3] N. Jing, *Quantum Kac-Moody algebras and vertex representations*, Lett. Math. Phys. **44** (1998), 261-271.

[M] I. G. Macdonald, *Symmetric functions and Hall polynomials*, 2nd ed., Clarendon Press, Oxford, 1995.

[Mc] J. McKay, *Graphs, singularities and finite groups*, Proc. Sympos. Pure Math. **37**, AMS (1980), 183–186.

[Se] G. Segal, *Unitary representations of some infinite dimensional groups*, Commun. Math. Phys. **80** (1980), 301–342.

[Sa] Y. Saito, *Quantum Toroidal algebras and their vertex representations*, Publ. Res. Inst. Math. Sci. **34** (1998), 155-177.

[VV] M. Varagnolo and E. Vasserot, *Double-loop algebras and the Fock space*, Invent. Math. **133** (1998), 133-159.

[W] W. Wang, *Equivariant K-theory and wreath products*, MPI preprint # **86**, August 1998; *Equivariant K-theory, wreath products and Heisenberg algebra*, math.QA/9907166, to appear in Duke Math. J.

[Z] A. Zelevinsky, *Representations of finite classical groups, A Hopf algebra approach.* Lecture Notes in Mathematics, **869**. Springer-Verlag, Berlin-New York, 1981.

Frenkel: Department of Mathematics, Yale University, New Haven, CT 06520

Jing: Department of Mathematics, North Carolina State University, Raleigh, NC 27695-8205 Mathematical Sciences Research Institute, 1000 Centennial Drive, Berkeley, CA 94720
E-mail address: `jing@math.ncsu.edu`

Wang: Department of Mathematics, North Carolina State University, Raleigh, NC 27695-8205 Department of Mathematics, Yale University, New Haven, CT 06520
E-mail address: `wqwang@math.ncsu.edu`